

Scotland's Rural College

Injurious tail biting in pigs: How can it be controlled in existing systems without tail docking?

D'Eath, RB; Arnott, G; Turner, SP; Jensen, T; Lahrmann, HP; Busch, ME; Niemi, JK; Lawrence, AB; Sandoe, P

Published in:
Animal

DOI:
[10.1017/S1751731114001359](https://doi.org/10.1017/S1751731114001359)

Print publication: 01/01/2014

Document Version
Peer reviewed version

[Link to publication](#)

Citation for pulished version (APA):

D'Eath, RB., Arnott, G., Turner, SP., Jensen, T., Lahrmann, HP., Busch, ME., Niemi, JK., Lawrence, AB., & Sandoe, P. (2014). Injurious tail biting in pigs: How can it be controlled in existing systems without tail docking? *Animal*, 8(9), 1479 - 1497. <https://doi.org/10.1017/S1751731114001359>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Injurious tail biting in pigs: how can it be controlled in existing systems
without tail docking?**

R.B. D'Eath¹, G. Arnott^{1,2}, S.P. Turner¹, T. Jensen³, H.P. Lahrmann³, M.E. Busch³,
J.K. Niemi⁴, A.B. Lawrence¹, P. Sandøe⁵

¹*SRUC, West Mains Road, Edinburgh, EH9 3JG, UK*

²*Queen's University Belfast, School of Biological Sciences, 97 Lisburn Road, Belfast,
BT9 7BL*

³*Danish Agriculture & Food Council, Pig Research Centre, Axeltorv 3, 1609
Copenhagen V, Denmark*

⁴*MTT Agrifood Research Finland, Economic Research, Kampusranta 9, FI-60320
Seinäjoki, Finland*

⁵*Department of Large Animal Sciences & Department of Food and Resource
Economics, University of Copenhagen, Grønnegårdsvej 8, 1870 Frederiksberg C,
Copenhagen, Denmark*

Corresponding author: Rick D'Eath. E-mail: rick.death@sruc.ac.uk

Running head: Controlling tail biting without tail docking

Abstract

Tail biting is a serious animal welfare and economic problem in pig production. Tail docking, which reduces but does not eliminate tail-biting, remains widespread. However, in the EU tail docking may not be used routinely, and some "alternative"

forms of pig production and certain countries do not allow tail docking at all. Against this background, using a novel approach focussing on research where tail injuries were quantified, we review the measures that can be used to control tail biting in pigs without tail docking. Using this strict criterion, there was good evidence that manipulable substrates and feeder space affect damaging tail biting. Only epidemiological evidence was available for effects of temperature and season, and the effect of stocking density was unclear. Studies suggest that group size has little effect, and the effects of nutrition, disease and breed require further investigation. The review identifies a number of knowledge gaps and promising avenues for future research into prevention and mitigation. We illustrate the diversity of hypotheses concerning how different proposed risk factors might increase tail biting through their effect on each other or on the proposed underlying processes of tail biting. A quantitative comparison of the efficacy of different methods of provision of manipulable materials, and a review of current practices in countries and assurance schemes where tail docking is banned, both suggest that daily provision of small quantities of destructible, manipulable natural materials can be of considerable benefit. Further comparative research is needed into materials, such as ropes, which are compatible with slatted floors. Also, materials which double as fuel for anaerobic digesters could be utilised. As well as optimising housing and management to reduce risk, it is important to detect and treat tail biting as soon as it occurs. Early warning signs before the first bloody tails appear, such as pigs holding their tails tucked under, could in future be automatically detected using precision livestock farming methods enabling earlier reaction and prevention of tail damage. However, there is a lack of scientific studies on how best to respond to outbreaks: the effectiveness of e.g. removing biters and/or bitten pigs, increasing enrichment, or applying

substances to tails should be investigated. Finally, some breeding companies are exploring options for reducing the genetic propensity to tail bite. If these various approaches to reduce tail biting are implemented we propose that the need for tail-docking will be reduced.

Keywords: Pigs, housing, enrichment, tail biting, behaviour

Implications

Tail biting in growing pigs is a serious welfare and economic problem, and there is pressure to avoid tail docking. For the first time relying only on studies where tail damage was recorded, we review the evidence on controlling tail biting in pigs that are not tail docked. Adequate feeder space and manipulable substrate provision are important, but more work is needed on the type and quantity of substrate needed. Vigilance for behavioural signs which occur before the first damaging biting would enable rapid detection and prevention/early response to outbreaks. Genetic selection could play a role in reducing tail biting.

Introduction

Tail biting in domestic pigs occurs when pigs bite and chew the tails of pen-mates. It is a considerable animal welfare (Munsterhjelm *et al.*, 2013) and economic problem, causing painful injuries which are a site for further infection (Sihvo *et al.*, 2012), resulting in carcass losses for producers (Kritas and Morrison, 2007; Valros *et al.*, 2004) and reducing weight gain (Sinisalo *et al.*, 2012; Wallgren and Lindahl, 1996). Several risk factors have been proposed, suggesting multi-factorial causation (EFSA, 2007; Schrøder-Petersen and Simonsen, 2001) and three different aetiologies have

75 been proposed (Taylor *et al.*, 2010). Removal of part of the tail (tail docking) a few
76 days after birth usually reduces the likelihood and severity of tail biting(Sutherland
77 and Tucker, 2011). Where tail docking is banned, tail biting incidence usually
78 increases, even when the housing environment and management are
79 improved(D'Eath *et al.*, 2014).

80
81 However, even though tail-docking reduces tail biting, it does not eliminate it and has
82 significant drawbacks: it is an acutely painful mutilation, and it may 'mask' the real
83 underlying problems in housing and management that result in tail biting(Sutherland
84 and Tucker, 2011). For these reasons, the EU Council Directive (2001/93/EC
85 amending Directive 91/630/EEC, The Council of The European Union, 2001b) came
86 into force from January 2003 banning the 'routine' tail-docking of pigs, unless 'there
87 is evidence that injuries ... to other pigs' ears or tails have occurred' and insisting that
88 before resorting to tail docking 'other measures shall be taken to prevent tail
89 biting...taking into account environment and stocking densities'. It goes on to state
90 that '...pigs must have permanent access to a sufficient quantity of material to
91 enable proper investigation and manipulation activities, such as straw, hay, wood,
92 sawdust, mushroom compost, peat or a mixture of such, which does not compromise
93 the health of the animals.' Despite this clear legal signal, tail-docking continues for
94 95% or more of pigs in European pig producing countries such asGermany,
95 Denmark, Belgium, France, Ireland, Netherlands and Spain, and for over 80% in the
96 UK (EFSA, 2007; Harley *et al.*, 2012).

97
98 Perhaps in response to this gap between policy and reality, the European
99 Commission (Directorate-General for Health and Consumers, DG Sanco; Bergersen,

2013) is currently engaging in a process to agree and clarify the guidance to farmers associated with the mentioned Directive and its later versions (the latest being 2008/120/EC on the protection of pigs). Some countries already go further than the EU directives in restricting tail docking (Mul *et al.*, 2010). In Denmark, no more than half of the tail may be docked, and in the Netherlands, a voluntary agreement exists between farmers and government to phase out tail docking entirely by 2023 (Spoolder *et al.*, 2011). A few countries already have either a complete ban on tail-docking (Sweden, Finland, Switzerland; EFSA, 2007; Swiss Federal Council, 2008) or a ban on docking without anaesthesia (Norway; EFSA, 2007) so that tail docking is rare. At the same time animal welfare protection organisations in many European countries focus on tail docking as a sign of welfare problems in intensive pig production; and in some countries political pressure is building up in favour of an effective ban on tail docking.

In another article, we consider the decisions facing farmers under current EU rules as to whether to tail dock, and the economic, legal and pig welfare consequences of this decision (D'Eath *et al.*, 2014). In the present article, we ask how farmers can become better at controlling tail biting without the use of tail docking. Our review focuses on changes that would be possible in existing systems, rather than considering radical system re-design (De Greef *et al.*, 2011). Various knowledge gaps are identified and promising areas for future innovation are proposed. We begin by introducing the nature of tail biting, and then review risk factors relating to the pigs' environment. For the first time, we rely only on studies which reported effects on tail injuries, rather than those which describe pigs' non-injurious interactions with tails ('tail-in mouth'). The relationships between these risk factors and the underlying

process(es) that govern the expression of tail biting are poorly understood, and we present a new illustration of the diversity of hypotheses. A novel illustrated meta-analysis quantifies the effectiveness of enrichment on tail biting in undocked pigs, and the practical experiences of countries and production systems in which tail docking is banned are considered. The next section of the review then focuses on risk factors that relate to characteristics of the pigs themselves, including the possibility of genetic selection to reduce tail biting. Finally we consider the prospects for early detection of tail biting outbreaks, possibly by automated means, which could facilitate targeted prevention measures. Farmers react to tail biting in various ways but little is known about the efficacy of these measures in preventing the further escalation of an outbreak.

Tail biting- why it remains an intractable problem

Tail biting occurs in outbreaks

Damaging tail-biting occurs in a sporadic way, in unpredictable 'outbreaks', rather like an infectious disease (Blackshaw, 1981). For example, in one study using abattoir data, 'high incidence farms' were identified at one point in time, but when a similar 'high incidence' list was made a few months later, most of the farms were different - although there were a few farms with a persistent problem (Busch *et al.*, 2004). In general, while some of the risk factors that affect the overall incidence of tail biting are known, for any given outbreak, the specific triggering factor(s) are usually difficult to identify. Sometimes a change (weather, season, food, or disease outbreak) can be identified, but often no obvious change has occurred, and the cause may be down to variability in individual pigs' threshold of response to risk factors.

Tail biting can spread quickly within the group

Tail biting begins with one pig in the pen starting to bite. Tail damage can increase rapidly, with one study reporting that progress from bite marks to a clearly visible tail wound took on average 7 days (Zonderland *et al.*, 2010b), although practical experience suggests that it can occur even more quickly. Over time, biting pigs may continue or escalate their biting of existing victims, but also begin biting other pigs in the group (Niemi *et al.*, 2011). Additionally, other pigs in an affected pen begin tail-biting too, perhaps because they copy the behaviour (social facilitation; Blackshaw, 1981) or the bitten tails might stimulate investigation and biting (stimulus enhancement; Fraser, 1987a). Although never formally studied, there appears to be considerable variation in the rate at which a pig increases its tail biting behaviour, and in the rate of spread to new biters. In one study, a batch of pigs already showing tail biting, moved to an environment with considerable space and access to rooting substrates, subsequently showed healing and improvement over time (De Greef *et al.*, 2011), suggesting that escalation is not inevitable.

Scientific investigation of tail biting is difficult

Tail biting is challenging to study. Its apparently sudden, unpredictable appearance and rapid spread can make it hard to investigate the events immediately before and after an outbreak begins. Its sporadic occurrence also means that a number of experimental studies have failed to observe any damaging tail biting at all. Such studies often report the effects of experimental treatments on tail investigation behaviour ('tail in mouth' Petersen *et al.*, 1995; Schroder-Petersen *et al.*, 2004), which is at best an indirect indicator of tail biting, because 'tail in mouth' behaviour may or may not be a pre-cursor to damaging tail biting (EFSA, 2007). Other studies include

all pig-directed oral behaviours including ear and flank biting, and sometimes belly-nosing, together in a single category(e.g. Jensen *et al.*, 2010; Zwicker *et al.*, 2013). This lack of precision makes interpretation difficult if the focus is on tail biting alone(Taylor *et al.*, 2010). In order to avoid these problems with indirect or imprecise indicators, this review focuses on studies where tail damage (with evidence of partial tail loss or of injury severe enough that blood was drawn) was the end point.

The sporadic occurrence of tail biting, and difficulties with experimental studies mean that multi-farm epidemiological studies (Goossens *et al.*, 2008; Moinard *et al.*, 2003)or abattoir data (Harley *et al.*, 2012; Valros *et al.*, 2004), sometimes combined with farm surveys (Hunter *et al.*, 2001), are often used to study tail biting. These usually record tail damage, and can find risk factors associated with it, but unlike experiments, are unable to determine cause and effect, so must be interpreted with caution. We begin by looking at risk factors in the pigs' environment, and then explore risk factors intrinsic to the pig. Some of these risk factors and causes of tail biting may also affect the related problems of ear- and flank-biting (Brunberg *et al.*, 2011), but this is beyond the scope of this paper.

Risk factors for tail biting in the pigs' environment and how to manage them

Tail biting does not have a single cause. It is a multi-factorial problem, and a variety of risk factors have been identified which are associated with it. Various efforts have been made to review all the currently known risk factors to weight their importance in order to influence policy makers (Bracke *et al.*, 2006; EFSA, 2007; Spoolder *et al.*, 2011) and to provide practical advice to farmers (Bracke *et al.*, 2004; Jensen *et al.*, 2004; Taylor *et al.*, 2012).

Taylor *et al*(2010) in a recent review made a convincing case that there were at least two and possibly three different types of tail biting: two-stage, sudden-forceful and obsessive. 'Two stage' tail biting results from re-directed foraging due to a lack of suitable substrates. There is a progression from investigation and gentle manipulation of tails (stage 1) to damaging biting (stage 2). The second type, 'sudden forceful' tail biting is an aggressive behaviour(Moinard *et al.*, 2003; Van Putten, 1969)apparently resulting from frustration over a lack of access to food, water or lying space. Pigs approaching a fully occupied resource such as a feeder may resort to biting at tails as the most readily available target for aggression. For example, in a recent study, 60% of the tail biting by pigs, which had limited feeder access (Palander *et al.*, 2013), occurred within 1m of the feeder (A. Valros pers. comm.). The third type, 'obsessive tail biting' is characterised by certain individual pigs which appear to be fixated on tails and go from one tail to another, inflicting damaging bites (Beattie *et al.*, 2005; Van de Weerd *et al.*, 2005). Here, we consider 'obsessive' biters to be individuals which are more likely than other pigs either to begin or to continue tail biting through the mechanisms explained above (and illustrated in Figure 1) for two-stage or sudden forceful tail biting.

The mechanism of action of each possible risk factor on the underlying processes controlling the expression of tail biting is in many cases unknown. Figure 1 illustrates many of the possible connections between proposed environmental risk factors and the underlying processes of 'two stage' and 'sudden-forceful' tail-biting (Taylor *et al.*, 2010). The nature of each possible connection is described in Supplementary Material S1. Evidence for the effect of risk factors on damaging tail biting is discussed further below.

Availability of manipulable materials

Manipulable materials which are attractive to pigs as measured by their motivation to access them (Holm *et al.*, 2008; Jensen *et al.*, 2008) or by the time pigs spend interacting with them over a sustained period have the characteristics 'ingestible', 'odorous', 'chewable', 'deformable' and 'destructible' (Studnitz *et al.*, 2007; Van de Weerd *et al.*, 2003; Van de Weerd and Day, 2009). The opportunity to perform investigation and manipulation behaviours are in themselves important for pig welfare (Studnitz *et al.*, 2007; Van de Weerd and Day, 2009), but here we focus on whether manipulable materials can reduce damaging tail biting apparently by providing an alternative outlet for investigatory behaviour.

Difficulties with the provision of loose manipulable materials on the floor

Systems making use of full or part-slatted floors, enabling automatic collection of pig faeces and urine (slurry) are common in indoor pig production. In comparison with straw-bedded systems, the labour (cleaning and waste handling) and input costs (e.g. of straw, peat and other substrates) are lower (Bornett *et al.*, 2003), some environmental impacts may be lower (Stern *et al.*, 2005), and liquid slurry is more valuable as a fertilizer than solid manure (Sanchez and Gonzalez, 2005). The requirement to provide manipulable materials to occupy pigs, presents a difficulty for farmers with systems which rely on slatted-floors and liquid slurry handling (via pumps). Materials such as long (unchopped) straw do not easily pass through slats leading to pen fouling. Additionally, too much straw can separate from the liquid slurry and build up in the slurry pit, or if it does flow, it can block parts of slurry-handling systems, such as holes, pipes or vacuum-based slurry pumps (Day *et al.*, 2008; Tuytens, 2005). Ways to reduce the problem of straw blockage may include

use of chopped straw, in combination with engineering solutions, such as larger diameter pipes (Evira, 2013), slurry pumps fitted with chopper blades, the use of smaller, faster flowing slurry systems (PRC, 2011), or progressive cavity pumps which are suitable for viscous liquids. Depending on the quantity and type of substrate, these measures may not be 100% effective but are more likely to be successful if considered at the building design stage.

Non-destructible materials such as metal chains, or rubber or hard plastic objects have been tried. Although pigs may initially interact with these due to their novelty, interest in them usually declines rapidly over a few days (Van de Perre *et al.*, 2011; Van de Weerd *et al.*, 2003). Even a repeating cycle of different objects may not be enough, as re-introduction of the same object after an interval of several weeks is usually not as effective at sustaining interest as a novel object would be (Van de Perre *et al.*, 2011). The European Commission have made it clear that chains and other non-destructible materials are not sufficient to comply with the EU Council directive (EC, 2009). Gradually destructible materials, which take days or weeks to be chewed through such as wooden poles (often mounted vertically in a tube at the side of the pen, or suspended from a chain) are popular with farmers in some countries as they require less regular replenishment than other more readily destructible substrates and appear to comply with the EU Council directive (2001/93/EC) which lists wood as a suitable material. Wooden poles were found to be an effective enrichment for reducing tail damage in a recent unpublished Finnish study using freshly felled tree trunks 5-10cm in diameter suspended on chains horizontally below snout level (Telkänranta *et al.*, 2014). However, since some of the features required to make a material attractive to pigs are lacking (ingestible,

odorous) or weak (chewable, deformable, destructible) in hard wood poles, the need to use soft, fresh woods which do have these features could be important.

In the face of these difficulties, an important question is whether it is possible to provide sufficient manipulable materials to pigs within existing intensive housing systems in order to reduce tail-biting to a level which is acceptably low from a management, production and welfare perspective without the need to tail dock (D'Eath *et al.*, 2014).

Alternative ways of providing manipulable materials

In part-slatted floored pens, it is possible to provide loose material such as chopped straw, peat or sawdust, which in small quantities may be used with slurry pumps (Munsterhjelm *et al.*, 2009). Substrate can be provided on the solid floor, while pigs defecate and urinate in the slatted part. To limit the passage of substrate from the solid to the slatted part of the floor, pen designs incorporating barriers (e.g. 50 mm high wooden strip, Zwicker *et al.*, 2013) or where the slatted area is raised (BPEX, 2010) may be used. Practical experience suggests that such designs are usually not entirely successful, especially in high temperatures where pigs may choose to defecate in the lying area, wallowing in the wet faeces to keep cool, and at higher stocking densities where functional separation of lying and dunging areas becomes more difficult to achieve, particularly in older pigs (Jensen *et al.*, 2012).

Faecal contamination of manipulable substrates is a common problem which reduces their attractiveness to pigs (Scott *et al.*, 2009), and this contamination can be reduced by hanging objects in the pen. Hanging of substrates limits the form of

interaction, for example chewing may be possible but not rooting (Day *et al.*, 2008). This might be important, or different forms of investigatory behaviour may substitute for one another in preventing tail biting, as long as the pigs are occupied. Hanging objects thus may have potential: in a meta-analysis of the time spent by pigs interacting with enrichment, properties promoting this interaction included enrichments which were suspended and/or deformable (Averós *et al.*, 2010). For example pigs show sustained interest in interacting with destructible ropes (Trickett *et al.*, 2009), or hanging objects with an edible component (Van de Weerd *et al.*, 2003), and 'flavoured rope' devices for pigs are being sold commercially in Finland. However, the effects of these forms of enrichment on tail biting have not been investigated.

Another approach is to deliver loose manipulable materials by means of an elevated rack, so that pigs can gradually obtain the material for themselves over a period (Beattie *et al.*, 2001; Van de Weerd *et al.*, 2006; Zwicker *et al.*, 2012; Zwicker *et al.*, 2013). This has the potential advantage of 'double' interaction (in the rack, and beneath it: on the floor, or in a box or feeder; Zwicker *et al.*, 2012) which might mean less material can be used for the same total amount of interest from the pigs. A related approach is to use a low-level rooting box which can contain loose materials and keeps them separate from slats (De Greef *et al.*, 2011; Van de Weerd *et al.*, 2003).

Quantifying the effects of different enrichment methods on tail damage

Studies published in refereed journals which compare the effect of different types and quantities of manipulable substrates on tail damage are summarised in Table

1. Most of the studies had pigs with intact tails, but some were docked, as indicated in the table legend. The studies all focus on grower-finisher pigs, except for Zonderland *et al.* (2008) which used weaners. Different indices of tail damage were used by different authors: Most studies report either the percentage of pigs removed from the study with severe tail injury, or the percentage of pigs or of pens having tail wounds. One study (Munsterhjelm *et al.*, 2009) used a tail lesion index (scoring from 0 to 2). To compare studies that used different measures, we calculated the fold-change in tail damage for each pair of treatments in these studies (i.e. the reduction in tail injury following the provision of one type of manipulable substrate compared to another). Where one of a pair of substrates had zero damage, it was not possible to calculate a fold-change, so 'max' was reported in Table 1, and this value did not contribute to the mean fold change, probably resulting in an underestimate of the effect size. Most studies compared deep straw with either no enrichment, or with minimal enrichment with chains or hanging toys, considered to represent commercial practice. In Figure 2, the information from the studies in Table 1 is summarised graphically, giving a quantification of the relative value of different materials as it is drawn to scale using the mean log 'fold' difference between observed levels of biting damage as the distance between the materials. A log scale was used so that 'fold' differences could be added together on the same scale and shown relative to each other in a single diagram (since e.g. $\log 2 + \log 3 = \log 6$).

In terms of manipulable substrate treatments that are compatible with fully or partially slatted floors, straw racks and light straw ($\leq 20\text{g/pig/day}$) are probably the most promising treatments for which data are available. Provision of straw in racks reduces tail damage compared to a rubber hose, chain or hanging toy, with two

studies finding a small but consistent reduction in the percentage of pens with tail wounds (fold-improvement of 1.9 or 1.7, Van de Weerd *et al.*, 2006; Zonderland *et al.*, 2008). In one study, the straw rack affected minor tail injuries, but was more effective at reducing severe tail damage (Van de Weerd *et al.*, 2006), which suggests that the straw rack might have reduced the rate of escalation of biting.

Light straw (10g twice a day per pig, Zonderland *et al.*, 2008), or light chopped straw and wood shavings (12.5g a day per pig, Munsterhjelm *et al.*, 2009) were both highly effective at reducing tail damage compared to minimally enriched treatments in two studies, with fold-differences almost as high as for deep straw studies (see Figure 2). Unfortunately, neither of these studies included plentiful loose material as a positive control. In a producer survey combined with an abattoir study, Hunter *et al.* (2001) found that 'light straw' use reduced tail biting damage risk compared to no straw. Despite these positive findings, chopped straw may not be as attractive to pigs as long straw: A behaviour study comparing chopped with long straw (each 400g/pig/day) suggested that chopped straw offers fewer possibilities for interaction and observed tail-biting behaviour increased (Day *et al.*, 2008). However, this study included non-injurious chewing and biting, and tail damage was not reported. In contrast, a Danish study suggested that chopped and long straw (each at 100g/pig/day) occupied pigs for a similar amount of time (Lahrmann and Steinmetz, 2011).

Our comparative survey has identified a number of data gaps: Only two studies included comparisons of more than one pair of treatments, allowing the substrates to be placed into an overall ranking (Van de Weerd *et al.*, 2006; Zonderland *et al.*,

2008), and there was a paucity of studies investigating how different quantities of straw or other materials affect tail damage. Time spent exploring and manipulating straw rather than other pigs increases with straw quantity until above 300g/pig/day (Olsson, 2011) or at around 500g/pig/day (Pedersen *et al.*, 2013). However, tail biting occurred at very low levels in these studies, even in treatments with only 20g/pig/day (undocked pigs, Olsson, 2011) or 10g/pig/day (docked pigs, Pedersen *et al.*, 2013). Also, no studies have compared hanging toys with no enrichment, and none have looked at the effect of hanging destructible enrichments such as ropes on tail damage, except for one recent report in suckling piglets (Telkänranta *et al.*, 2014b), so there is considerable scope for further research.

Manipulable materials as fuel for anaerobic digesters

Materials which act as foraging enrichment for pigs could double as fuel for anaerobic digesters (AD). This idea is being tested in the “Starplus” system at Wageningen (Verdoes, 2014). ADs enable farmers to deal with farm wastes, producing energy (methane) and digestates which can be used as fertiliser. Pig slurry provides micronutrients and trace elements needed for bacterial growth, but its energy content is low, so (non-wood) biological materials are added, some which could provide rooting/foraging (and eating) opportunities for pigs: chopped grass, maize or grass silage, sugar beet and kitchen waste (if concerns over biosecurity could be addressed). For example, pigs prefer chopped straw mixed with Maize silage over straw (Jensen *et al.*, 2010; Jensen and Pedersen, 2007) possibly because it may include edible components. Many questions remain, however: materials must be compatible with floor slats/slurry systems, a method to deliver substrate to the pens is required, fungal growth in wet fermenting materials can be a

problem (T. Jensen pers. comm.), and there are hygiene issues if pigs are eating material from the floor. Finally, fuel source costs, energy prices and government policies affect the economic feasibility of AD.

Social factors- space allowance (stocking density), group size, mixing

At space allowances lower than those currently recommended in the EU, reduced space allowance increased tail damage in one experiment (Krider *et al.*, 1975). At space allowances closer to or within the recommended range, one multi-farm study found an association between reduced space allowance and tail injuries (Goossens *et al.*, 2008) but another similar study did not (Smulders *et al.*, 2008), and no effect was found in an experimental study (Street and Gonyou, 2008). Group size (Schmolke *et al.*, 2003; Smulders *et al.*, 2008; Street and Gonyou, 2008) and mixing of groups (Smulders *et al.*, 2008; Zonderland *et al.*, 2008) had no effect in studies where tail damage was reported.

Feeding- feeder space, feed restriction, feed type, nutrients, minerals

Restricted feeder space increased tail biting in one experimental study (damaged tails, Hansen *et al.*, 1982) and is a risk factor in epidemiological studies (Hunter *et al.*, 2001; Moinard *et al.*, 2003). Other experimental studies, in which low levels of tail biting occurred, found no effect of feeder space (Georgsson and Svendsen, 2001; Georgsson and Svendsen, 2002). The form and presentation of feed may be important: pigs fed pelleted diets showed higher levels of tail injury than meal or liquid fed pigs in one study (Hunter *et al.*, 2001) while Temple *et al.* (2012) found liquid feed in a trough increased tail injury compared to wet feed in a hopper.

Nutritional qualities of the diet: protein, specific amino acids, minerals or high energy density have all been suggested to affect tail biting(Edwards, 2011), but there is little direct evidence of nutritional manipulations affecting tail damage. Experiments using 'model' tails suggest that attraction to blood may be increased if the diet is nutritionally inadequate in terms of protein (Fraser *et al.*, 1991) or minerals (Fraser, 1987b).Tail biting pigs were more attracted to cords soaked with pig blood than their non-biting pen mates(McIntyre and Edwards, 2002b) and this preference can be reduced by the addition of the amino acid tryptophan to their diets (McIntyre and Edwards, 2002a). Differences in serotonin metabolism in the prefrontal cortex and an altered pattern of tryptophan uptake have been reported in tail biting pigs in contrast to bitten and unaffected group-mates or unaffected pigs from another group (Valros *et al.*, 2013). Additional salt in the diet or on the floor of the pen can increase foraging and drinking behaviour(Brooks, 2005), which may reduce biting, but it is not clear whether this was effectively a foraging enrichment or addressing a nutritional deficiency. Jaeger *et al.*(2013)proposed a novel causal pathway for tail biting: high energy density diets for weaner pigs (as well as exposure to various pathogens) result in a build-up of endotoxins which cause ear or tail necrosis, which then attracts biting. If the necrotic tissue is itchy, this could increase the tolerance of tail investigation and biting in victim pigs.

Climate- temperature, draughts, seasonal effects

Either low (Temple *et al.*, 2012), or both low and high temperatures (Geers *et al.*, 1989) have been identified by epidemiological studies as risk factors for tail damage, and providing access to a water misting system can reduce tail injury in hot climates (Courboulay *et al.*, 2008). Seasonal effects on tail damage have been identified

(Schröder-Petersen and Simonsen, 2001, Busch pers. comm.), the exact nature of which varies between different studies. It seems plausible that rapid changes in temperature (either up or down), an increase in draughts at certain times of year (known to affect activity, Scheepens *et al.*, 1991), or heat stress are likely to be the underlying cause of seasonal effects (Figure 1), as there is a limit to the capacity of ventilation/heating/cooling systems in most pig buildings.

Disease, including parasitism

Disease has been proposed to be a risk factor (Edwards, 2011). Levels of tail damage are higher in herds with higher levels of respiratory illness (Elst *et al.*, 1988 cited by Edwards 2011; Moinard *et al.*, 2003), and in a study where health records from individual pigs were examined, leg disorders and tail damage were highly correlated (Niemi *et al.*, 2012). Caution is required with the interpretation of epidemiological data, as disease may result from infections that follow tail biting (Kritas and Morrison, 2007; Moinard *et al.*, 2003), or poor health status may be an indirect indicator of less technically efficient farms.

Controlled studies in which measures to improve health result in a reduction in tail damage provide better evidence. Currently there is only an anecdotal report of tail biting being reduced following anthelmintic treatment (Barnikol, 1978) and an as-yet unpublished study concerning PCV2 vaccination (Parker *et al.* in prep, cited by Edwards, 2011). So at this stage, the evidence for disease as a cause of tail biting is weak.

The experience of countries and assurance schemes where tail docking is banned

Pig producers in some countries and assurance schemes have already had to adapt their systems to cope with greater restrictions on tail docking, and the changes they have made are instructive. The tail docking and housing system rules for grower-finisher pigs applied by selected non-docking European countries and selected assurance schemes are summarised in Table 2, with some systems which permit tail docking included for comparison.

The 'tail docking restricted or banned' farms have a number of features in common, many of which may reduce tail biting risk. The space allowance is usually more generous, with up to 50% more space per pig being provided. Fully slatted pens are not allowed, enabling manipulable materials to be provided on the solid-floored part of the pen (although partially slatted, part drained floors are permitted in Finland). Compared to the EU minimum provision, there are more specific rules on the quantity of materials, usually by specifying the frequency of replenishment, or the behaviour that pigs must be able to perform: in Finland pigs must be able to make small piles of material, Freedom Food requires sufficient quantities of material for rooting, pawing and chewing behaviour. The type of material provided is often also restricted, for example Sweden and the Danish assurance scheme Antonius require straw, and Norway repeats the EU list, but stipulates wood chips rather than 'wood' which rules out the use of wooden posts.

The small scale of farms in Finland, Norway and Switzerland, compared to the UK and Denmark (Table 2) might enable a greater supervision of the animals (assuming

more staff per pig), making detection and prevention of tail biting easier, and smaller farms often have lower disease risk (Goldberg et al 2000). For example, Finland is free from Porcine Reproductive and Respiratory disease, and mycoplasma and Salmonella are at low levels, although a lower density of farms and fewer pig movements including imports may also be important here.

Tail docking is not completely outlawed in all of the assurance schemes in Table 2. The assurance schemes Antonius, Outdoor (including Organic) in Denmark and Freedom Food allow farmers to apply for a dispensation to use tail docking for a limited time if a tail biting outbreak occurs. For example, Freedom Food farmers must annually seek written permission to dock, and if tail biting in the previous year was low, they are encouraged to trial a cessation of docking for some pigs, with the aim of stopping docking altogether. At each application, farmers must document the other measures they have taken to prevent tail biting and quantify their success. The standards give a detailed list of a number of environmental improvements that should be tried including providing straw and increasing feeder space. In 2010, 30% of Freedom Food breeding farms supplying indoor wean to finish herds requested permission to dock (Kate Parkes RSPCA pers. comm.). This suggests that the majority of scheme members are managing to rear intact tailed pigs, while those with tail biting problems are allowed to use taildocking to protect the welfare of potential victim pigs.

Finally, Table 2 gives figures on tail biting prevalence in the different countries and schemes, estimated from abattoirs. It is very difficult to compare the figures, since they are not collected in a standardised way, for example pigs with missing tails are

usually not counted as injured(EFSA, 2007; Keeling *et al.*, 2012). It is also difficult to compare between docked and undocked pigs; if the number of lesions are counted, long tails provide a greater area for biting than docked or part-docked tails (Webster and Day, 1998). To avoid this problem, it might be best to use data on partial carcass condemnations (PCC), which can be used as indicators of the most severe cases of tail biting(Kritas and Morrison, 2007; Valros *et al.*, 2004), but these are not always available. Occasionally, different housing systems are assessed in the same way during the same period at one abattoir. Data from a Danish study(Forkman *et al.*, 2010) show that tail biting damage at slaughter was higher in intact-tailed pigs in organic (or outdoor)systems (range 1.0 – 4.0%), in comparison to docked pigs in conventional indoor housing (range 0.5-1.5%; Table 2). These data suggest that tail docking is more effective at reducing tail biting than the combined effect of various improvements to the environment(as reported by Hunter *et al.*, 2001). It also highlights how challenging it is for producers to rear intact-tailed pigs, even in improved environments. However it would be better to have data comparing countries and systems which used a standard scoring method and included PCC.

Risk factors for tail biting which are characteristics of the pig and how to manage them

Characteristics of the pigs themselves can affect their propensity to tail bite or to be bitten. This could involve any step of the tail biting process (Figure 1). To take the ‘two-stage’ form of tail biting for example pigs could vary in:i) how much they perform exploratory behaviour(e.g. Zonderland *et al.*, 2011), ii) whether they explore tails rather than other things, iii) whether tail manipulation becomes biting, iv) whether one bite becomes many, and v) in the likelihood of learning tail biting from a pen

mate. Also, differences between pigs in their propensity to tail bite or be bitten could occur due to a greater sensitivity to any environmental risk factor (Figure 1).

Characteristics of victim pigs: sex and breed

Certain pigs may be more likely to become victims of tail biting. In several studies, castrated males were more likely to be the victims of tail biting than females (Kritas and Morrison, 2004; Kritas and Morrison, 2007; tail damage, Wallgren and Lindahl, 1996), with the risk to males rising with the proportion of females in the pen (Kritas and Morrison, 2004). Tail biting was higher in pigs grouped by sex (intact males and females) in one abattoir study (Hunter *et al.*, 2001), but no relationship was found in an on-farm study of castrates and females (Steinmetz and Pedersen, 2009). All-female pens have been reported as having more (Zonderland *et al.*, 2010a) tail damage than pens of entire males. In another study, all female pens had less (Steinmetz and Pedersen, 2009) tail damage than pens of all-castrated males. It is not clear what causes these different findings in relation to sex.

Breed may affect the likelihood of being bitten. In a Swedish pedigree population of male pigs with low levels of tail biting, bitten pigs were: Yorkshire (3.5%), Landrace (1.8%) and Hampshire (0.1%, Westin, 2003). However this was based on small numbers (63 victims out of 3049) in mixed breed groups which were not composed in a systematic way. Thus the evidence for breed differences in the risk of being bitten is weak.

Characteristics of tail biters: growth retardation, early experience, breed

It has been suggested that tail biters are often the smaller pigs in a group (Sambraus, 1985) which is in agreement with data from some studies (Zonderland *et al.*, 2011) although others have found that tail biters were no more likely to be smaller than average (Breuer *et al.*, 2005). Once tail biting begins, certain individual pigs show much higher levels of biting than others (Beattie *et al.*, 2005; Van de Weerd *et al.*, 2005), and have been characterised as ‘fanatical’ or ‘obsessive’ biters (Taylor *et al.*, 2010). In one study with small numbers of ‘obsessive’ biters, these pigs were smaller than average (Van de Weerd *et al.*, 2005). Beattie *et al.* (2005) found that ‘tail-in-mouth’ behaviour was higher in pigs which grew poorly in the first 3 weeks after weaning. Although poorly supported by evidence from studies which include tail damage, it remains a popular theory with farmers, that smaller pigs in a pen begin biting perhaps because they resort to this biting as an aggressive tactic when excluded from food (Schrøder-Petersen and Simonsen, 2001) and/or because of a problem with nutrition or metabolism (Edwards, 2006; EFSA, 2007).

EFSA (2007) concluded that the rearing environment is not as important as the current environment for tail-biting risk. Where pigs receive manipulable materials during the grower stage, past housing experience makes little difference to tail manipulation behaviours (Day *et al.*, 2002; no tail damage reported, Simonsen, 1995; Statham *et al.*, 2011). However, a greater risk of tail lesions caused by tail biting occurs in pigs which have experienced manipulable materials in the farrowing pen early in life, but which are then absent during later stages (Munsterhjelm *et al.*, 2009; Ruiterkamp, 1985). In contrast to the small effects seen in experimental studies, epidemiological studies have found associations between early life factors and tail biting. These factors include slatted floors (Smulders *et al.*, 2008) or absence of

substrates(Moinard *et al.*, 2003) in the farrowing pen, and limited feeder space or high temperatures in the nursery (Smulders *et al.*, 2008). Epidemiological studies of course do not prove a causal link and further research is required.

As well as breed differences in the propensity to become victims of tail biting (described above), some have reported breed differences in the propensity to perform tail biting. In the Swedish pedigree population with mixed breed pens of male pigs described earlier, the biters were: Landrace (1.7%), Yorkshire (0.64%) and Hampshires (0.1%), but only 27 biters out of 3049 animals were observed(Westin, 2003).In a UK study, there was no effect of breed on tail biting, although there were breed differences in ear biting (Duroc>Large White>Landrace; Breuer *et al.*, 2003).Two other studies reported finding no breed differences in the performance of tail biting(Guy *et al.*, 2002; Lund and Simonsen, 2000).Thus, as with breed differences in being tail bitten, breed differences in bitingmight occur, but the evidence is fairly weak.

Genetics of tail biting: biters, victims and unaffected pigs

A single published quantitative genetic study exists which found that biting other pigs' tails was a heritable trait (Breuer *et al.*, 2005), at least in Landrace (but not in Large White) pigs. Heritability was low at 0.05 ± 0.02 , although tail-biters were rare (295 tail biters in a population of 9018 pigs) and tail biting was treated as a binary trait, which reduces the power of genetic analysis.

Commercial pig breeding mainly focuses on economically important traits of lean growth rate, food conversion and reproductive traits such as litter size. Some pig

breeding companies are considering broadening their breeding goals, and traits relating to behaviour and welfare issues such as tail biting are of interest (Canario *et al.*, 2013; Merks *et al.*, 2012). The inclusion of additional traits in a breeding index inevitably leads to a reduced rate of genetic progress in other traits (Falconer and Mackay, 1996). However, breeding companies normally use economic weightings in breeding indices, and the considerable costs of tail biting could make it economically optimal to include a trait linked to lowered levels of tail biting in a multi-trait index (Lawrence *et al.*, 2004)..

A number of other factors stand in the way of conventional genetic selection against tail biting (see discussion in Turner, 2011). The positive genetic correlation relationship between tail biting and lean tissue growth rate found by Breuer *et al.* (2005) could slow genetic progress if found in other populations. Phenotyping is also a challenge: identification of 'biting' pigs is considerably more difficult than identification of victims. Direct observation of biting may be the most accurate method, but is time-consuming, especially since tail biting, as described above, often occurs in sporadic, unpredictable outbreaks. Also, it may be important to identify the individual pig which starts the outbreak ('first biter'), as once bloody tails appear in the pen, other pigs are more likely to begin biting. Because of these difficulties, there would be enormous value in identification of a proxy trait, associated with tailbiting. Unfortunately, tests based on artificial tails have proved largely disappointing in terms of their predictive value for real tail biting (Beattie *et al.*, 2005; Breuer *et al.*, 2003; Statham, 2008), although in one study the time spent manipulating an enrichment device before tail biting began was higher in biters than victims (Zonderland *et al.*, 2011). Automated detection of tailbiting might be possible

using similar methods to those proposed for detection of early-warning signs (see next section).

There is some prospect of identifying molecular genetic markers of pigs at lower risk of tail biting, an approach which reduces the amount of phenotyping required. Single Nucleotide Polymorphism (SNP) markers of biting and victim pigs (in contrast to non-biting controls from the same pen) have been identified (Wilson *et al.*, 2012). Brain gene expression studies also suggest that biters and victims have more in common than unaffected pigs from the same group (Brunberg *et al.*, 2013a) or a different group (Brunberg *et al.*, 2013b). These authors suggest that unaffected pigs may show a 'tail-biting resistant' phenotype. If confirmed in other populations, this idea suggests that selection against both biters and victims and for unaffected pigs might be possible.

Another approach which side-steps the problem of phenotyping tail biters is to use 'associative genetic effects' (Camerlink *et al.*, 2012; Turner, 2011). Quantitative genetic models for pig growth can be modified allowing pigs to have heritable influences on the growth of their pen-mates (Bijma *et al.*, 2007a; Bijma *et al.*, 2007b; Rodenburg *et al.*, 2010). Depending on the context, these 'social breeding values' might reflect differences in positive social behaviours such as social nosing (Camerlink *et al.*, 2012) or in negative behaviours such as aggression, food competition, disease transmission and ear, flank or tailbiting. To have a more direct effect on tail biting, modelling of associative genetic effects could be used in combination with phenotyping for tail damage. With sufficient representation of

different sires across pens, pigs with a high genetic propensity to cause tail damage to pen mates could be identified, without the need to observe biting behaviour.

Selection to reduce tail biting behaviour could raise ethical concerns, particularly concerning 'naturalness' (D'Eath *et al.*, 2010). Because 'two-stage' tail-biting results from frustrated foraging behaviour, we could speculate that selection for lower tail biting might also reduce foraging. Also, selecting animals to function well in poor environments, rather than improving the animals' environment to satisfy their needs might seem distasteful to some, and could lead to a decline in housing standards (Kanis *et al.*, 2004). However, given that pigs are already undergoing constant genetic change to alter production traits, alongside improvements to the housing environment we should perhaps consider whether genetic selection to reduce tail biting could be part of a solution which makes an end to tail docking possible (D'Eath *et al.*, 2014).

Although this is speculative, breeding to reduce tail length might be possible as tail length is heritable in various mammalian species (rodents, Barnett, 1965; sheep, Branford Oltenacu and Boylan, 1974; cats, Howell and Siegel, 1966), and naturally short-tailed pigs might be less prone to becoming victims of tail-biting. There are probably difficulties though as tail length is likely to be genetically correlated with back length (a desired trait in bacon pigs) and tail-less mutations may have undesirable side-effects such as those seen in Manx cats (Howell and Siegel, 1966). Even if breeding to reduce tail length were successful, other concerns remain. The curly tail could be seen by consumers as an essential pig characteristic (although it is absent in wild pig species) and may have a function in

communication (Kiley-Worthington, 1976). Finally, breeding rather than docking to shorten tails still sidesteps the problem that pigs need an outlet for their foraging behaviour.

Early detection and targeted prevention

An alternative approach to the problem of tail-biting is to detect outbreaks before or as soon as they begin, and to carry out targeted intervention (such as those discussed in the next section) to ameliorate or even prevent an outbreak (FAWC, 2011). Regardless of the system, if pig producers could identify certain 'at risk' individuals, groups, or batches, and target them for preventive intervention, this would be cheaper and more practical compared to making changes for every pig.

Early detection of tail biting might be possible by identifying changes in pig behaviour that precede an outbreak. Four main types of early warning sign have been described, which appear in the days or weeks before an outbreak begins (first bloody tails): 1) General activity ('restlessness') increases (Statham *et al.*, 2009; Zonderland *et al.*, 2011), particularly in biters (Svendsen *et al.*, 2006). 2) Non-damaging 'tail in mouth' behaviour increases (Feddes and Fraser, 1994; Fraser, 1987a; Schrøder-Petersen and Simonsen, 2001). 3) Tails are held down or 'tucked under' (Statham *et al.*, 2009; Zonderland *et al.*, 2009; 2010). 4) Feeding patterns might change. In one study, feeder visits tended ($p < 0.1$) to be lower in groups which went on to tail bite 6-9 weeks pre-outbreak, and tended ($p < 0.1$) to increase during weeks 2-5 pre-outbreak in pigs which would become tail biting victims (Wallenbeck and Keeling, 2013). More research on feeding patterns is needed.

Increased observation of pigs by staff might identify these early warning signs, but staff time has a cost, so automatic detection ('precision livestock farming') would be attractive (reviewed by Rushen *et al.*, 2012). The detection of specific behaviours such as tail posture and tail in mouth behaviour may be possible (Sonoda *et al.*, 2013), but increases in activity (and perhaps feeding patterns) are perhaps the easiest of the 'early warning signs' to detect automatically (Costa *et al.*, 2013). One promising approach is 'optical flow', which estimates animal activity by quantifying overall pixel changes from moment to moment in a video image. 'Optical flow' has been used to detect the reduced activity of lame broiler chickens (Dawkins *et al.*, 2009), and the disturbance of behaviour in laying hen flocks when feather pecking is occurring (Lee *et al.*, 2011).

The use of on-board animal devices (such as electronic ID ear tags), combined with detectors in the pen to record pig location also has potential to detect changes in activity (or feeding patterns). Currently, the infrastructure and consumable costs associated with either video or EID approaches to monitoring pig behaviour may be prohibitive. But with falling costs and various other benefits of electronic ID (easier record keeping for medicines and at weighing, or even for detecting when a pig has not visited the feeder in a long while) and video (estimation of pig size average and variability) the use of these technologies could become more widespread in future. Detection of changes in pig vocalisations is also a plausible approach (Manteuffel *et al.*, 2004) but would require considerable further research and validation.

Reacting to outbreaks

Once a tail-biting outbreak occurs, pig producers react in various ways (Arey, 1991). Hunter *et al.* (2001) surveyed British pig producers and found that 67% removed the

bitten pig(s), 51% added enrichment objects, 25% applied sprays or tar to injured tails, 16% added straw, 6% reduced stocking density and 6% gave antibiotics. In one study, moving pigs that were already biting to pens with substrates and more space resulted in reduced biting behaviour (De Greef *et al.*, 2011), and Edwards (2011) has suggested that the effectiveness of salt or other nutritional supplements should be investigated. There is considerable scope for more research in this area: Only one scientific study of the effectiveness of interventions has been reported. Zonderland *et al* (2008) compared the interventions of removing the biter or adding straw as soon as tail damage was detected and found them to be equally effective (the 'no intervention' control was considered unethical).

Where pigs are removed from a tail biting group, bitten pigs are easier to identify (Hunter *et al.*, 2001), but removing tail biters might have a greater impact. Since biting spreads rapidly to other pigs, the time window is small for removing the first biter. Even if biters are removed, leaving bitten pigs in the pen might encourage new biters (because they are attracted to the bloody or scabbed tails), so it has been suggested that removal of *both* biters *and* bitten pigs might be optimal (Boyle and Lemos Teixeira, 2010; Zonderland *et al.*, 2008).

The other difficulty with removing pigs is the question of how to manage the pigs that are removed (Boyle and Lemos Teixeira, 2010). In an outbreak where multiple pigs are removed, the farmer may be constrained by space to group house them. They must also decide whether to use a lower stocking density and/or substrates or other forms of enrichment for those removed pigs. There is obviously a concern that removing biters and putting them into new groups could result in further tailbiting,

although anecdotally, Zonderland *et al* (2008) reported that they did not experience this problem. Removing pigs to different groups and/or returning them needs to be carefully managed as it can result in social aggression including fighting and bullying (Marchant-Forde and Marchant-Forde, 2005).

Another intervention worthy of further research is the application of aversive substances to tails. Bracke (2009) found that when pigs were offered untreated ropes, or ropes treated with Dippel's oil or Stockholm tar to chew on, they avoided the treated ropes, suggesting that these treatments might be aversive when applied to tails. On the other hand, a concern over adding substances (including antibiotic sprays) to tails is that it might make them more novel, stimulating investigation and perhaps biting.

As suggested by Edwards (2011), there is clearly an urgent need for systematic research into the effectiveness of different methods for reacting to outbreaks. This research should: i) investigate the different methods separately or in combination and develop others, ii) investigate the quantity/frequency of enrichment that is necessary to reduce tail biting iii) investigate the optimal timing of interventions (there may be a point after which certain methods cease to be effective), iv) investigate whether it is most effective and efficient to target individuals, pens, or a whole room of pigs.

Conclusions

The risk factors affecting tail damage caused by tail biting were reviewed. A number of risk factors that have been proposed and reviewed elsewhere (EFSA, 2007; Schrøder-Petersen and Simonsen, 2001) are not currently well supported by

experimental studies where damaging tail biting was the end point. These include group size, nutrition, disease incidence and pig breed. Surprisingly, the evidence for an effect of stocking density was also quite weak. Epidemiological evidence alone suggests that temperature and season might be important. The evidence was strongest for the provision of manipulable substrates, and an effect of feeder space was also found.

Housing systems using slatted floors and liquid slurry handling are in widespread use due to their economic advantages, but limit the amount of loose manipulable substrates that can be used. A crucial question for this review was whether, at commercial stocking densities, in part solid-, partly slatted-floored pens, small quantities of straw or similar manipulable substrate (perhaps delivered via a rack), can reduce tail-biting to the point where tail docking is no longer necessary. Very few studies have looked at this, but the few that have were promising. Damaging tail biting was greatly reduced in two studies with undocked pigs using light straw (10g twice a day per pig, Zonderland *et al.*, 2008) or light chopped straw and wood shavings (12.5g a day per pig, Munsterhjelm *et al.*, 2009), and the experience of Finland which uses small quantities of enrichment material is also positive. Further studies investigating the effect of quantity and type of enrichment material on tail biting risk are necessary, and such studies are especially valuable if treatments are compared to a negative control of very little enrichment and a positive control of a plentiful loose material. In particular, further studies of destructible hanging materials such as ropes and destructible fresh wood would be useful. As well as controlled scientific studies, investigations into the experiences of producers in assurance schemes which are working to phase out tail docking would also be worthwhile. As a

way of reducing the cost of using enrichment materials, the possibility of using materials which could combine with pig slurry as fuel for anaerobic digesters is interesting but faces a number of technical hurdles.

The mechanisms by which the various proposed environmental risk factors might affect the underlying process(es) of tail biting are largely unknown. Possible mechanisms are shown in Figure 1, but much of this is speculative (see Supplementary Material S1) and there is considerable scope for further research into whether and how these risk factors might cause or affect tail biting. Alongside optimising the environment, it may be possible to use genetic selection to reduce tail biting. The challenge of phenotyping by identifying biters (Breuer *et al.*, 2005) and especially the ‘first biter’ could potentially be made easier (using automatic detection or proxy measures), made into a smaller task (by identification of genetic markers) or side-stepped altogether (by the use of ‘associative genetic effects’ for growth, or possibly, for tail lesions). The possibility that a ‘tail-biting resistant’ phenotype might exist is interesting (Brunberg *et al.*, 2013b), but identifying these pigs would still be challenging; they are the pigs in an affected pen which are neither biters nor victims.

Another potential area for innovation is the use of precision livestock farming methods to automatically detect the early warning signs of a tail biting outbreak at the pre-damaging stage. Various behavioural signs have been identified, including tail position, ‘tail in mouth’ behaviour, and increased activity, some of which might be detectable by automatic methods based on electronic tags (see <http://pigit.ku.dk> and www.pigwise.eu) or on video (Sonoda *et al.*, 2013). If farmers could identify when and where an outbreak of tail-biting was about to begin, they could target preventative

measures, which would be more economic in terms of time and materials than making changes for all pigs. A final potential area for innovation is to test the efficacy of measures to stop tail-biting once it begins (or just before it begins), which has been the subject of only one scientific study (Zonderland *et al.*, 2008).

Spoolder *et al.*(2011) suggested that ‘an intact curly tail can be regarded as the single most important welfare indicator in finishing pigs, since to achieve this requires a high standard of housing and management over a pig’s lifetime, so it serves as an ‘iceberg indicator’ of welfare (FAWC, 2009) and demonstrates respect for the ‘animal integrity’ of the pig. Within a system type, it also indicates good management to prevent (or quickly deal with)tail biting.A current difficulty is that alternative systems with intact-tailed pigs usually suffer from higher levels of tailbiting than conventional systems that tail dock(Hunter *et al.*, 2001; Table 2). This means there is an ethical question as to how we should weigh a welfare impact on many (all pigs being docked as a precaution) with a worse welfare impact for a few (victims of tail biting)(D'Eath *et al.*, 2014). Would we consider that a ban on tail docking had led to improved welfare if it increased tail damage at slaughter from 1% of pigs to 4% of pigs?The threshold for what constitutes an‘acceptably low level’of tail biting must be decided in a wider ethical debate which considers the pigs’ perspective. The experience of countries with complete bans on tail docking is that farmers do learn to reduce tail biting in other ways, although the resulting economic costs of this adaptation may reduce competitiveness and participation in export markets.

In most EU countries where docking is permitted, the letter of the EU Directive (2001/93/EC) that requires provision of manipulable materials is being followed,

although the pressure group Compassion in World Farming found that on the farms they visited in a number of EU countries it was not (CIWF, 2008). However, the Directive states that docking should be used as a last resort only when there is evidence of a tail biting problem and other environmental measures have been tried, and this is only being enacted in reality by a minority of producers, for example some assurance schemes (e.g. Danish Antonius and Organic, UK Freedom Food). This approach seems a logical middle road (which appears to be the 'spirit' of the EU directive), allowing the majority of pigs in these schemes to benefit by avoiding docking and having their behavioural needs met, while still allowing docking to protect the welfare of potential tail biting victims on farms with a problem. However, it would be more difficult to enforce than a complete ban on tail docking. Thus it would require considerably more detail in terms of the measures producers should take before resorting to tail docking, and these measures would most likely involve substantial changes from current practice, imposing considerable costs on producers.

Acknowledgements

Funding from the Danish Pig Research Centre supported this review. SRUC receives funding from the Rural and Environment Science and Analytical Services (RESAS) Division of the Scottish Government. Bouda Ahmadi, Emma Baxter, Sarah Ison, Karsten Klint Jensen and Kenny Rutherford contributed to valuable discussions and provided comments on drafts. Roland Weber, Linda Keeling, Kate Parkes and Lene Juul Pedersen also provided valuable information.

References

895 Arey DS 1991. Tail-Biting in Pigs. *Farm Building Progress* 20-23.

896 Averós X, Brossard L, Dourmad JY, De Greef KH, Edge HL, Edwards SA and Meunier-
897 Salaün MC 2010. A meta-analysis of the combined effect of housing and environmental
898 enrichment characteristics on the behaviour and performance of pigs. *Applied Animal*
899 *Behaviour Science* 127, 73-85.

900 Barnett SA 1965. Genotype and environment in tail length in mice. *Experimental Physiology*
901 50, 417-429.

902 Barnikol M 1978. Observations of tail biting (cannibalism) amongst fattening hogs and
903 breeding pigs in connection with protein and mineral nourishment. *Tieraerztliche Umschau*
904 33, 540.

905 Beattie VE, Breuer K, O'Connell NE, Sneddon IA, Mercer JT, Rance KA, Sutcliffe MEM and
906 Edwards SA 2005. Factors identifying pigs predisposed to tail biting. *Animal Science* 80,
907 307-312.

908 Beattie VE, Sneddon IA, Walker N and Weatherup RN 2001. Environmental enrichment of
909 intensive pig housing using spent mushroom compost. *Animal Science* 72, 35-42.

910 Bergersen B 2013. Extended working group meeting on the development of guidelines
911 concerning Council Directive 2008/120/EC on the protection of pigs. European Commission
912 Directorate-General for Health and Consumers (DG-Sanco). Brussels, Belgium.

913 Bijma P, Muir WA and van Arendonk JAM 2007a. Multilevel selection 1: Quantitative
914 genetics of inheritance and response to selection. *Genetics* 175, 277-288.

915 Bijma P, Muir WM, Ellen ED, Wolf JB and van Arendonk JAM 2007b. Multilevel selection 2:
916 Estimating the genetic parameters determining inheritance and response to selection.
917 *Genetics* 175, 289-299.

918 Blackshaw JK 1981. Some behavioural deviations in weaned domestic pigs: persistent
919 inguinal nose thrusting, and tail and ear biting. *Animal Production* 33, 325-332.

920 Bornett HLI, Guy JH and Cain PJ 2003. Impact of Animal Welfare on Costs and Viability of
921 Pig Production in the UK. *Journal of Agricultural & Environmental Ethics* 16, 163-186.

922 Boyle L and Lemos Teixeira D 2010. Improving pig welfare will reduce carcass losses.
923 http://www.teagasc.ie/publications/view_publication.aspx?PublicationID=48 (retrieved
924 21/10/2013). Teagasc Pig Farmers Conferences. Moorepark, Ireland, pg 47-56.

925 BPEX 2010. Environmental enrichment for pigs (retrieved on 21/10/2013).
926 <http://www.bpex.org.uk/articles/301028>,

927 Bracke MBM 2009. Rope test may indicate efficacy of tail-biting treatments in growing pigs.
928 *Animal Welfare* 18, 263-266.

929 Bracke MBM, Hulsegge B, Keeling L and Blokhuis HJ 2004. Decision support system with
930 semantic model to assess the risk of tail biting in pigs. 2. 'Validation'. *Applied Animal*
931 *Behaviour Science* 87, 45-54.

932 Bracke MBM, Zonderland JJ, Lenskens P, Schouten WGP, Vermeer H, Spoolder HAM,
933 Hendriks HJM and Hopster H 2006. Formalised review of environmental enrichment for pigs
934 in relation to political decision making. *Applied Animal Behaviour Science* 98, 165-182.

935 Branford Oltenacu EA and Boylan WJ 1974. Inheritance of tail length in crossbred
936 Finnsheep. *Journal of Heredity* 65, 331-334.

937 Breuer K, Sutcliffe MEM, Mercer JT, Rance KA, Beattie VE, Sneddon IA and Edwards SA
938 2003. The effect of breed on the development of adverse social behaviours in pigs. *Applied*
939 *Animal Behaviour Science* 84, 59-74.

940 Breuer K, Sutcliffe MEM, Mercer JT, Rance KA, O'Connell NE, Sneddon IA and Edwards SA
941 2005. Heritability of clinical tail-biting and its relation to performance traits. *Livestock*
942 *Production Science* 93, 87-94.

943 Brooks, P. H. 2005. Effect of diet on the behaviour and welfare of pigs *Proceedings of the*
944 *Manitoba Swine Seminar* retrieved on 21-10-2013 from [http://www.prairieswine.com/effect-](http://www.prairieswine.com/effect-of-diet-on-the-behaviour-and-welfare-of-pigs/)
945 [of-diet-on-the-behaviour-and-welfare-of-pigs/](http://www.prairieswine.com/effect-of-diet-on-the-behaviour-and-welfare-of-pigs/).

946 Brunberg E, Jensen P, Isaksson A and Keeling LJ 2013a. Brain gene expression differences
947 are associated with abnormal tail biting behavior in pigs. *Genes, brain, and behavior* 12,
948 275-281.

949 Brunberg E, Wallenbeck A and Keeling LJ 2011. Tail biting in fattening pigs: Associations
950 between frequency of tail biting and other abnormal behaviours. *Applied Animal Behaviour*
951 *Science* 133, 18-25.

952 Brunberg E, Jensen P, Isaksson A and Keeling LJ 2013b. Behavioural and Brain Gene
953 Expression Profiling in Pigs during Tail Biting Outbreaks- Evidence of a Tail Biting Resistant
954 Phenotype. *PLoS ONE* 8, e66513.

955 Busch, ME, Wachmann, H, Nielsen, EO, and Petersen, HH 2004. Halebid - Kan slagtedata
956 anvendes til at identificere besætninger med høj forekomst? (Tail biting - can slaughter data
957 be used to identify herds with a high prevalence?). Danish Pig Research Centre (PRC),
958 Copenhagen, Denmark. Report: Meddelelse 643,
959 http://vsp.lf.dk/Publikationer/Kilder/lu_medd/2004/643.aspx?full=1 (downloaded on
960 21/10/2013).

961 Camerlink I, Bijma P, Kemp B and Bolhuis JE 2012. Relationship between growth rate and
962 oral manipulation, social nosing, and aggression in finishing pigs. *Applied Animal Behaviour*
963 *Science* 142, 11-17.

964 Canario L, Mignon-Grasteau S, Dupont-Nivet M and Phocas F 2013. Genetics of
965 behavioural adaptation of livestock to farming conditions. *Animal* 7, 357-377.

966 CIWF 2008. The State of Europe's Pigs: An exposé. *Compassion in World Farming*,
967 Investigative Report,

968 CIWF 2009. Welfare of Pigs in the European Union- The urgent need for reform of existing
969 legislation and effective enforcement. *Compassion in World Farming*, Godalming, Surrey,
970 UK.

971 Costa A, Ismayilova G, Borgonovo F, Leroy T, Berckmans D and Guarino M 2013. The use
972 of image analysis as a new approach to assess behaviour classification in a pig barn. *Acta*
973 *Veterinaria Brno* 82, 25-30.

974 Council of State 2002. Valtioneuvoston asetus eläinsuojeluasetuksen muuttamisesta. VNa
975 910/2002 (Government decree on the regulation of the protection of animals) retrieved on
976 21-10-2013 from <http://www.finlex.fi/fi/laki/alkup/2002/20020910>.

977 Council of State 2012. Valtioneuvoston asetus sikojen suojelusta. VNa 629/2012
 978 (Government decree on the protection of pigs) retrieved on 21-10-2013 from
 979 <http://www.finlex.fi/fi/laki/alkup/2012/20120629>.

980 Courboulay V, Eugene A and Delarue E 2009. Welfare assessment in 82 pig farms: effect of
 981 animal age and floor type on behaviour and injuries in fattening pigs. *Animal Welfare* 18,
 982 515-521.

983 Courboulay V, Souffran H, Guingand N, Massabie P and Dubois A 2008. Intérêt de la
 984 brumisation en engraissement pour améliorer le confort des porcs en croissance (Fogging system: a
 985 way to improve the welfare of growing-finishing pigs during summer periods). *Journées*
 986 *Recherche Porcine* 40, 243-246.

987 D'Eath RB, Niemi J, Ahmadi B, Arnott G, Turner SP, Baxter EM, Rutherford KMD, Jensen T,
 988 Lahrmann HP, Busch ME, Jensen KK, Lawrence AB and Sandøe P 2014. Tail docking vs.
 989 tail biting- a scientific, legal, socio-economic and ethical analysis. *Animal Welfare* submitted.

990 D'Eath RB, Conington J, Lawrence AB, Olsson IAS and Sandøe P 2010. Breeding for
 991 behavioural change in farm animals: Practical, economic and ethical considerations. *Animal*
 992 *Welfare* 19(S), 17-27.

993 Danish Crown 2007. Kvalitetshåndbog for Antonius griseproduktion (Quality handbook for
 994 Antonius pig production) Copenhagen, Denmark retrieved on 21-10-2013 from
 995 <http://www.danishcrown.dk/lib/file.aspx?fileID=286>.

996 Danish Government 2000. Lov no. 104 af 14 Februar 2000 om indendørs hold af smågrise,
 997 avls- og slagtesvin (Act on indoor keeping of pigs, breeding and fattening). Copenhagen,
 998 Denmark.

999 Danish Government 2003a. BEK nr 323 af 06/05/2003, Bekendtgørelse om beskyttelse af
 1000 svin (Order on the protection of pigs) Fødevareministeriet (Danish Ministry of Food)
 1001 Copenhagen, Denmark retrieved on 21-10-2013a from
 1002 <https://www.retsinformation.dk/Forms/R0710.aspx?id=1571>.

1003 Danish Government 2003b. BEK nr 324 af 06/05/2003, Bekendtgørelse om halekupering og
 1004 kastration af dyr (Order on tail docking and castration of animals) Fødevareministeriet
 1005 (Ministry of Food) Copenhagen, Denmark retrieved on 21-10-2013b from
 1006 <https://www.retsinformation.dk/Forms/R0710.aspx?id=1572>.

1007 Dawkins MS, Lee HJ, Waitt CD and Roberts SJ 2009. Optical flow patterns in broiler chicken
 1008 flocks as automated measures of behaviour and gait. *Applied Animal Behaviour Science*
 1009 119, 203-209.

1010 Day JEL, Burfoot A, Docking CM, Whittaker X, Spooler HAM and Edwards SA 2002. The
 1011 effects of prior experience of straw and the level of straw provision on the behaviour of
 1012 growing pigs. *Applied Animal Behaviour Science* 76, 189-202.

1013 Day JEL, De Weerd HAV and Edwards SA 2008. The effect of varying lengths of straw
 1014 bedding on the behaviour of growing pigs. *Applied Animal Behaviour Science* 109, 249-260.

1015 De Greef KH, Vermeer HM, Houwers HWJ and Bos AP 2011. Proof of Principle of the
 1016 Comfort Class concept in pigs. Experimenting in the midst of a stakeholder process on pig
 1017 welfare. *Livestock Science* 139, 172-185.

1018 Defra 2003. Code of recommendations for the welfare of livestock: pigs. Defra Publications,
1019 London.

1020 DVFA 2013. Fødevarestyrelsens vejledning om beskæftigelses- og rodematerialer (DVFA
1021 guidance on employment and enrichment materials) Danish Veterinary and Food
1022 Administration (Fødevarestyrelsen) Copenhagen, Denmark retrieved on 21-10-2013 from
1023 <http://www.foedevarestyrelsen.dk/Leksikon/Sider/Hold-af-svin.aspx>.

1024 EC 2009. Answer from the European Commission to a written question (E-5360/2009) from
1025 Member of the European Parliament Caroline Lucas Strasbourg retrieved on 21-10-2013
1026 from <http://www.europarl.europa.eu/sides/getAllAnswers.do?reference=E-2009-5360&language=EN>.
1027

1028 Edwards S 2011. What do we know about tail biting today? The Pig Journal 66, 81-86.

1029 Edwards SA 2006. Tail biting in pigs: Understanding the intractable problem. Veterinary
1030 Journal 171, 198-199.

1031 EFSA 2007. The risks associated with tail biting in pigs and possible means to reduce the
1032 need for tail docking considering the different housing and husbandry systems (Question No
1033 EFSA-Q-2006-013). Annex to the EFSA Journal 611, 1-13.

1034 Elst WET, Vaessen MA, Vos HJMP, Binnendijk GP, Huime REM and Backus GBC 1988.
1035 Proefverslag- Varkensproefbedrijf- Zuid-enwest-Nederland 200, 1-54.

1036 Eurostat 2013. Livestock: number of farms and heads of animals by type of farming in 2010
1037 European Commission retrieved on 21-10-2013 from
1038 <http://epp.eurostat.ec.europa.eu/portal/page/portal/agriculture/data/database>.

1039 Evira 2013. Sikojen tonkimismateriaali (Materials for pigs to root) Evira (Finnish Food Safety
1040 Authority) Helsinki, Finland retrieved on 21-10-2013 from
1041 http://www.evira.fi/files/attachments/fi/elaimet/elainsuojelu_ja_elainten_pito/sikojen_tonkimis_materiaali.pdf.
1042

1043 Falconer, D.S. & Mackay, T.F.C. 1996. Introduction to Quantitative Genetics. 4th edn. New
1044 York: Longman.

1045 FAWC 2009. Farm Animal Welfare in Great Britain: Past, Present and Future. Farm Animal
1046 Welfare Council, London.

1047 FAWC 2011. Opinion on mutilations and environmental enrichment in piglets and growing
1048 pigs. Farm Animal Welfare Council, London.

1049 Feddes JJR and Fraser D 1994. Non-nutritive chewing by pigs: Implications for tail-biting
1050 and behavioural enrichment. Transactions of the ASAE 37, 947-950.

1051 Fjetland O and Kjastad HP 2002. Tail biting in pigs. Norsk Veterinartidsskrift 114, 249-253.

1052 Forkman, B., Olsen, P., Nørgaard, N., Laursen, P. B., Pedersen, L. J., Holm, L., Viekilde, K.,
1053 and Søndergaard, J. T. 2010. Arbejdsgrupperapport om hold af svin (Working group report
1054 on the keeping of pigs) Justitsministeriet Departementet Copenhagen, Denmark retrieved on
1055 21-10-2013 from
1056 <https://bdkv2.borger.dk/Lovgivning/Hoeringsportalen/Sider/Fakta.aspx?hpid=2146002290>.

1057 Fraser D 1987a. Attraction to blood as a factor in tail-biting by pigs. *Applied Animal*
1058 *Behaviour Science* 17, 61-86.

1059 Fraser D 1987b. Mineral-deficient diets and the pig's attraction to blood: implications for tail-
1060 biting. *Canadian Journal of Animal Science* 67, 909-918.

1061 Fraser D, Bernon DE and Ball RO 1991. Enhanced attraction to blood by pigs with
1062 inadequate dietary-protein supplementation. *Canadian Journal of Animal Science* 71, 611-
1063 619.

1064 Friland 2012. Frilandgris Produktionskoncept - en god begyndelse (Outdoor pigs production
1065 concept- a good start) Randers, Denmark retrieved on 21-10-2013 from
1066 <http://www.friland.dk/lib/file.aspx?fileID=636>.

1067 Geers R, Dellaert B, Goedseels V, Hoogerbrugge A, Vranken E, Maes F and Berckmans D
1068 1989. An Assessment of Optimal Air Temperatures in Pig Houses by the Quantification of
1069 Behavioral and Health-Related Problems. *Animal Production* 48, 571-578.

1070 Georgsson L and Svendsen J 2001. One or two feeders for groups of 16 growing-finishing
1071 pigs: Effects on health and production. *Acta Agriculturae Scandinavica Section A-Animal*
1072 *Science* 51, 257-264.

1073 Georgsson L and Svendsen J 2002. Degree of competition at feeding differentially affects
1074 behavior and performance of group-housed growing-finishing pigs of different relative
1075 weights. *Journal of Animal Science* 80, 376-383.

1076 Goldberg, T.L., Weigel, R.M., Hahn, E.C. & Scherba, G. 2000. Associations between genetics,
1077 farm characteristics and clinical disease in field outbreaks of porcine reproductive and
1078 respiratory syndrome virus. *Preventive Veterinary Medicine* 43, 293-302.

1079 Goossens X, Sobry L, Odberg F, Tuytens F, Maes D, De Smet S, Nevens F, Opsomer G,
1080 Lommelen F and Geers R 2008. A population-based on-farm evaluation protocol for
1081 comparing the welfare of pigs between farms. *Animal Welfare* 17, 35-41.

1082 Guy JH, Rowlinson P, Chadwick JP and Ellis M 2002. Behaviour of two genotypes of
1083 growing/finishing pig in three different housing systems. *Applied Animal Behaviour Science*
1084 75, 193-206.

1085 Hansen LL, Hagelsø AM and Madsen A 1982. Behavioural results and performance of
1086 bacon pigs fed ad libitum from one or several self-feeders. *Applied Animal Ethology* 8, 307-
1087 333.

1088 Harley S, More S, O'Connell N, Hanlon A, Teixeira D and Boyle L 2012. Evaluating the
1089 prevalence of tail biting and carcass condemnations in slaughter pigs in the Republic and
1090 Northern Ireland, and the potential of abattoir meat inspection as a welfare surveillance tool.
1091 *Veterinary Record* 171, 621-+.

1092 Holm L, Jensen MB, Pedersen LJ and Ladewig J 2008. The importance of a food feedback
1093 in rooting materials for pigs measured by double demand curves with and without a common
1094 scaling factor. *Applied Animal Behaviour Science* 111, 68-84.

1095 Holmgren N and Lundeheim N 2004. Risk factors for tail biting. *Proceedings of the 18th*
1096 *IPVS congress. Hamburg, Germany*, pg 786.

- 1097 Howell JM and Siegel PB 1966. Morphological Effects of the Manx Factor in Cats. Journal of
1098 Heredity 57, 100-104.
- 1099 Hunter EJ, Jones TA, Guise HJ, Penny RHC and Hoste S 2001. The relationship between
1100 tail biting in pigs, docking procedure and other management practices. Veterinary Journal
1101 161, 72-79.
- 1102 Jaeger F 2013. Das Projekt „intakter Ringelschwanz“ beim Schwein - stehen wir vor dem
1103 Durchbruch? (The project "intact curly tails" in pigs - are we just before a breakthrough?).
1104 Tierärztliche Umschau 68, 3-11.
- 1105 Jensen MB and Pedersen LJ 2007. The value assigned to six different rooting materials by
1106 growing pigs. Applied Animal Behaviour Science 108, 31-44.
- 1107 Jensen MB, Studnitz M, Halekoh U, Pedersen LJ and Jorgensen E 2008. Pigs' preferences
1108 for rooting materials measured in a three-choice maze-test. Applied Animal Behaviour
1109 Science 112, 270-283.
- 1110 Jensen MB, Studnitz M and Pedersen LJ 2010. The effect of type of rooting material and
1111 space allowance on exploration and abnormal behaviour in growing pigs. Applied Animal
1112 Behaviour Science 123, 87-92.
- 1113 Jensen, T, Riis, AL, Maribo, H, Vernerisen, A, and Busch, ME 2004. Halebid Manual (Tail-
1114 biting manual). Videncenter for Svineproduktion og danske slagterier, Copenhagen,
1115 Denmark.
- 1116 Jensen T, Kold Nielsen C, Vinther J and D'Eath RB 2012. The effect of space allowance for
1117 finishing pigs on productivity and pen hygiene. Livestock Science 149[1]. pg 33-40.
- 1118 Jordbruksverket 2010. Statens jordbruksverks föreskrifter och allmänna råd om djurhållning
1119 inom lantbruket m.m. (Swedish State Board of Agriculture Regulations and Guidelines on
1120 Animal Husbandry in Agriculture etc.). Statens jordbruksverk, Jönköping, Sweden.
- 1121 Kanis E, van den Belt H, Groen AF, Schakel J and De Greef KH 2004. Breeding for
1122 improved welfare in pigs: a conceptual framework and its use in practice. Animal Science 78,
1123 315-329.
- 1124 Keeling LJ, Wallenbeck A, Larsen A and Holmgren N 2012. Scoring tail damage in pigs: an
1125 evaluation based on recordings at Swedish slaughterhouses. Acta veterinaria scandinavica
1126 54, 32.
- 1127 Kiley-Worthington M 1976. Tail Movements of Ungulates, Canids and Felids with Particular
1128 Reference to Their Causation and Function As Displays. Behaviour 56, 69-115.
- 1129 Krider JL, Albright JL, Plumlee MP, Conrad JH, Sinclair CL, Underwood L, Jones RG and
1130 Harrington RB 1975. Magnesium Supplementation, Space and Docking Effects on Swine
1131 Performance and Behavior. Journal of Animal Science 40, 1027-1033.
- 1132 Kritas SK and Morrison RB 2004. An observational study on tail biting in commercial grower-
1133 finisher barns. Journal of Swine Health and Production 12, 17-22.
- 1134 Kritas SK and Morrison RB 2007. Relationships between tail biting in pigs and disease
1135 lesions and condemnations at slaughter. Veterinary Record 160, 149-152.

1136 Lahrmann, HP and Steinmetz, H 2011. Effekt af halmlængde på forekomsten af unormal
1137 adfærd og halmrettet adfærd hos slagtesvin (Effect of straw length on the incidence of
1138 abnormal behavior and straw directed behavior of pigs). Danish Pig Research Centre (VSP),
1139 Copenhagen, Denmark.

1140 Lawrence AB, Conington J, and Simm G 2004 Breeding and animal welfare: practical and
1141 theoretical advantages of multi-trait selection. *Animal Welfare* 13, S191-S196.

1142 Lee HJ, Roberts SJ, Drake KA and Dawkins MS 2011. Prediction of feather damage in
1143 laying hens using optical flows and Markov models. *Journal of the Royal Society Interface* 8,
1144 489-499.

1145 LMD 2003. FOR 2003-02-18 nr 175: Forskrift om hold av svin (Regulations on the keeping of
1146 pigs) Landbruks- og matdepartementet (Norwegian Ministry of Agriculture and Food) retrieved on
1147 21-10-2013 from <http://www.lovddata.no/cgi-wift/ldles?doc=/sf/sf/sf-20030218-0175.html>.

1148 Lund A and Simonsen HB 2000. Aggression and stimulus-directed activities in two breeds of
1149 finishing pig. *The Pig Journal* 45, 123-130.

1150 MAF 1997. Sikojen pidolle asetettavat eläinsuojeluvaatimukset. (Animal protection
1151 requirements on concerning the keeping of pigs.). Decision 13/EEO/1997. Ministry of
1152 Agriculture and Forestry, Helsinki, Finland.

1153 Manteuffel G, Puppe B and Schon PC 2004. Vocalization of farm animals as a measure of
1154 welfare. *Applied Animal Behaviour Science* 88, 163-182.

1155 Marchant-Forde JN and Marchant-Forde RM 2005. Minimizing inter-pig aggression during
1156 mixing. *Pig News and Information* 26, 63N-73N.

1157 McIntyre J and Edwards SA 2002a. An investigation into the effect of tryptophan on tail chewing
1158 behaviour of growing pigs. *Proceedings of the British Society of Animal Science*. pg 34.

1159 McIntyre J and Edwards SA 2002b. Preference for blood and behavioural measurements of
1160 known tailbiting pigs compared to control penmates. *Proceedings of the 36th International
1161 Congress of the International Society for Applied Ethology*. Egmond an Zee, Netherlands, pg
1162 93.

1163 Merks JWM, Mathur PK and Knol EF 2012. New phenotypes for new breeding goals in pigs.
1164 *Animal* 6, 535-543.

1165 MFLF 2012. Vejledning om økologisk jordbrugs-produktion (Guidance on organic agricultural
1166 production) Ministeriet for Fødevarer, Landbrug og Fiskeri (Danish Ministry of Food,
1167 Agriculture and Fisheries) Copenhagen, Denmark retrieved on 21-10-2013 from
1168 http://1.naturerhverv.fvm.dk/oekologi_vejledning_bedrifter.aspx?ID=2137.

1169 Moinard C, Mendl M, Nicol CJ and Green LE 2003. A case control study of on-farm risk
1170 factors for tail biting in pigs. *Applied Animal Behaviour Science* 81, 333-355.

1171 Mul, M, Vermeij, I, Hindle, V, and Spoolder, H 2010. EU Welfare Legislation on Pigs. Report
1172 273 Livestock Research, Wageningen UR,

1173 Munsterhjelm C, Brunberg E, Heinonen M, Keeling L and Valros A 2013. Stress measures in
1174 tail biters and bitten pigs in a matched case-control study. *Animal Welfare* 22, 331-338.

1175 Munsterhjelm C, Peltoniemi OAT, Heinonen M, Halli O, Karhapää M and Valros A 2009.
 1176 Experience of moderate bedding affects behaviour of growing pigs. *Applied Animal*
 1177 *Behaviour Science* 118, 42-53.

1178 Niemi JK, Sinisalo A, Valros A and Heinonen M 2011. The timing and treatment of tail biting in
 1179 fattening pigs. Hulgren, J., Persson, P., Nadeau, E., and Fogelberg, F. 24th NJF Congress
 1180 "Food, Feed, Fuel and Fun- Nordic Light on Future Land Use and Rural Development", 14-
 1181 16th June, NJF Report 7. Uppsala, Sweden.

1182 Niemi JK, Sinisalo A, Valros A and Heinonen M 2012. Hännänpurenta - syy vai seuraus?
 1183 (Tail biting - cause or effect?). Schulman, N. and Kauppinen, H. *Maataloustieteen Päivät*
 1184 2012 [Verkkojulkaisu] (Agricultural Science Days 2012 [Online]). Suomen
 1185 Maataloustieteellisen Seuran julkaisu no. 28 (The Scientific Agricultural Society of Finland
 1186 publications no. 28). Helsinki, Finland (<http://www.smts.fi>).

1187 Olsson, L. 2011. Optimizing Amount of Straw for Growing-Finishing Pigs- Considering Time
 1188 Spent in Manipulative Behaviour. SLU, Uppsala, Sweden.

1189 Palander PA, Heinonen M, Simpura I, Edwards SA and Valros AE 2013. Jejunal morphology
 1190 and blood metabolites in tail biting, victim and control pigs. *Animal FirstView*, 1-9.

1191 Partanen, K., Niemi, J., Voutilainen, L. & Jukola, E. 2012. Vastuullisen tuotannon vaatimusten
 1192 täyttäminen näkyy lihasikalan tuloksessa (Responsible pig meat production). In:
 1193 *Maataloustieteen Päivät 2012 [Verkkojulkaisu]* (Agricultural Science Days 2012 [Online])
 1194 (Ed. by N. Schulman & H. Kauppinen) Helsinki, Finland (<http://www.smts.fi>).

1195 Pedersen LJ, Herskin M and Forkman B 2013. How much straw do slaughter pigs need to
 1196 meet the need for exploration. ISAE 47th Congress. Florianopolis, Brazil, pg 46.

1197 Petersen V, Simonsen HB and Lawson LG 1995. The effect of environmental stimulation on
 1198 the development of behaviour in pigs. *Applied Animal Behaviour Science* 45, 215-224.

1199 PRC 2011. Annual Report. Pig Research Centre, Danish Agriculture and Food Council,

1200 Rodenburg TB, Bijma P, Ellen ED, Bergsma R, de Vries S, Bolhuis JE, Kemp B and van
 1201 Arendonk JAM 2010. Breeding amiable animals? Improving farm animal welfare by including
 1202 social effects into the genetic model. *Animal Welfare* 19, 77-82.

1203 RSPCA 2012. Welfare Standards for Pigs. Freedom Food
 1204 <http://www.rspca.org.uk/sciencegroup/farmanimals/standards/pigs>.

1205 Ruiterkamp, W.A. 1985. Het Gedrag Van Mestvarkens in Relatie Tot Huisvesting (The
 1206 Behaviour of Fattening Pigs in Relation to Housing). PhD Thesis, University of Utrecht, The
 1207 Netherlands.

1208 Rushen J, Chapinal N and de Passille A 2012. Automated monitoring of behavioural-based
 1209 animal welfare indicators. *Animal Welfare* 21, 339-350.

1210 Sambras, H.H. 1985. Mouth-Based Anomalous Syndromes. In: *World Animal Science*, A5,
 1211 *Ethology of Farm Animals. A Comprehensive Study of the Behavioural Features of Common*
 1212 *Farm Animals* pp. 391-422.

1213 Sanchez M and Gonzalez JL 2005. The fertilizer value of pig slurry. I. Values depending on
 1214 the type of operation. *Bioresource Technology* 96, 1117-1123.

- 1215 Scheepens CJM, Hessing MJC, Laarakker E, Schouten WGP and Tielen MJM 1991.
1216 Influences of intermittent daily draft on the behaviour of weaned pigs. *Applied Animal*
1217 *Behaviour Science* 31, 69-82.
- 1218 Schmolke SA, Li YZ and Gonyou HW 2003. Effect of group size on performance of growing-
1219 finishing pigs. *Journal of Animal Science* 81, 874-878.
- 1220 Schroder-Petersen DL, Heiskanen T and Ersboll AK 2004. Tail-in-mouth behaviour in
1221 slaughter pigs, in relation to internal factors such as: Age, size, gender, and motivational
1222 background. *Acta Agriculturae Scandinavica Section A-Animal Science* 54, 159-166.
- 1223 Schrøder-Petersen DL and Simonsen HB 2001. Tail biting in pigs. *Veterinary Journal* 162,
1224 196-210.
- 1225 Scollo A, Di Martino G, Bonfanti L, Stefani AL, Schiavon E, Marangon S and Gottardo F
1226 2013. Tail docking and the rearing of heavy pigs: The role played by gender and the
1227 presence of straw in the control of tail biting. Blood parameters, behaviour and skin lesions.
1228 *Research in Veterinary Science* 95, 825-830.
- 1229 Scott K, Chennells DJ, Armstrong D, Taylor L, Gill BP and Edwards SA 2007. The welfare of
1230 finishing pigs under different housing and feeding systems: liquid versus dry feeding in fully-
1231 slatted and straw-based housing. *Animal Welfare* 16, 53-62.
- 1232 Scott K, Chennells DJ, Campbell FM, Hunt B, Armstrong D, Taylor L, Gill BP and Edwards
1233 SA 2006. The welfare of finishing pigs in two contrasting housing systems: Fully-slatted
1234 versus straw-bedded accommodation. *Livestock Science* 103, 104-115.
- 1235 Scott K, Taylor L, Gill BP and Edwards SA 2009. Influence of different types of
1236 environmental enrichment on the behaviour of finishing pigs in two different housing systems
1237 3. Hanging toy versus rootable toy of the same material. *Applied Animal Behaviour Science*
1238 116, 186-190.
- 1239 Sihvo H, Simola O, Munsterhjelm C and Syrja P 2012. Systemic Spread of Infection in Tail-
1240 Bitten Pigs. *Journal of Comparative Pathology* 146[1]. pg 73.
- 1241 Simonsen HB 1995. Effect of early rearing environment and tail docking on later behaviour
1242 and production in fattening pigs. *Acta Agriculturae Scandinavica Section A- Animal Science*
1243 45, 139-144.
- 1244 Sinisalo A, Niemi JK, Heinonen M and Valros A 2012. Tail biting and production
1245 performance in fattening pigs. *Livestock Science* 143, 220-225.
- 1246 SLU and LRF 2009. Swedish Pig Production (Pamphlet). Swedish University of Agricultural
1247 Sciences and Swedish Pig Producers Association. Uppsala, Sweden.
- 1248 Smulders D, Hautekiet V, Verbeke G and Geerst R 2008. Tail and ear biting lesions in pigs:
1249 an epidemiological study. *Animal Welfare* 17, 61-69.
- 1250 Soil Association 2012. Organic standards: farming and growing (revision 16.7) UK retrieved
1251 on 21-10-2013 from <http://www.soilassociation.org/organicstandards>.
- 1252 Sonoda LT, Fels M, Oczak M, Vranken E, Ismayilova G, Guarino M, Viazzi S, Bahr C,
1253 Berckmans D and Hartung J 2013. Tail Biting in pigs - Causes and management intervention
1254 strategies to reduce the behavioural disorder. A review. *Berliner und Munchener*
1255 *Tierärztliche Wochenschrift* 126, 104-112.

1256 Spoolder, H, Bracke, M, Mueller-Graf, C, and Edwards, S 2011. Preparatory work for the
 1257 future development of animal based measures for assessing the welfare of pig - Report 2:
 1258 Preparatory work for the future development of animal based measures for assessing the
 1259 welfare of weaned, growing and fattening pigs including aspects related to space allowance,
 1260 floor types, tail biting and need for tail docking. Technical report submitted to EFSA,

1261 Statham,P. 2008. Effects of Experience and Individuality on Tail Biting in Pigs. PhD Thesis,
 1262 University of Bristol.

1263 Statham P, Green L, Bichard M and Mendl M 2009. Predicting tail-biting from behaviour of
 1264 pigs prior to outbreaks. *Applied Animal Behaviour Science* 121, 157-164.

1265 Statham P, Green L and Mendl M 2011. A longitudinal study of the effects of providing straw
 1266 at different stages of life on tail-biting and other behaviour in commercially housed pigs.
 1267 *Applied Animal Behaviour Science* 134, 100-108.

1268 Steinmetz, HV and Pedersen, ML 2009. Kønsvis opstaldnings betydning for forekomsten af
 1269 halebid (Single sex housing- its importance for tailbiting). Report Meddelelse nr. 845
 1270 Videncenter for Svineproduktion (Danish Pig Research Centre),

1271 Stern S, Sonesson U, Gunnarsson S, Öborn I, Kumm KI and Nybrant T 2005. Sustainable
 1272 Development of Food Production: A Case Study on Scenarios for Pig Production. *AMBIO: A*
 1273 *Journal of the Human Environment* 34, 402-407.

1274 Street BR and Gonyou HW 2008. Effects of housing finishing pigs in two group sizes and at
 1275 two floor space allocations on production, health, behavior, and physiological variables.
 1276 *Journal of Animal Science* 86, 982-991.

1277 Studnitz M, Jensen MB and Pedersen LJ 2007. Why do pigs root and in what will they root?
 1278 A review on the exploratory behaviour of pigs in relation to environmental enrichment.
 1279 *Applied Animal Behaviour Science* 107, 183-197.

1280 Sutherland MA and Tucker CB 2011. The long and short of it: A review of tail docking in farm
 1281 animals. *Applied Animal Behaviour Science* 135, 179-191.

1282 Svendsen J, Olsson AC and Botermans J 2006. Data on tail biting in pigs. Proceedings of
 1283 the 19th IPVS congress. Copenhagen, Denmark, pg 613.

1284 Swiss Federal Council 2008. 455.1 Animal Welfare Ordinance (TSchV) retrieved on 21-10-
 1285 2013 from <http://www.bvet.admin.ch/themen/tierschutz/02443/index.html?lang=de>.

1286 Taylor NR, Main DCJ, Mendl M and Edwards SA 2010. Tail-biting: A new perspective. *The*
 1287 *Veterinary Journal* 186, 137-147.

1288 Taylor NR, Parker RMA, Mendl M, Edwards SA and Main DCJ 2012. Prevalence of risk
 1289 factors for tail biting on commercial farms and intervention strategies. *The Veterinary Journal*
 1290 194, 77-83.

1291 Telkänranta H, Bracke M and Valros A 2014a. Developing chewable and rootable low-cost
 1292 objects for commercial pig farms. Proceedings of the 25th ISAE Nordic Regional Meeting.
 1293 Drobak, Norway, pg 10.

1294 Telkänranta,H., Swan,K., Hirvonen,H. & Valros,A. 2014b. Chewable materials before
 1295 weaning reduce tail biting in growing pigs. *Applied Animal Behaviour Science* in press
 1296 (online at <http://dx.doi.org/10.1016/j.applanim.2014.01.004>)

1297 Temple D, Courboulay V, Velarde A, Dalmau A and Manteca X 2012. The welfare of growing
1298 pigs in five different production systems in France and Spain: assessment of health. *Animal*
1299 *Welfare* 21, 257-271.

1300 The Council of The European Union 2001a. Commission Directive 2001/93/EC of 9
1301 November 2001 amending Directive 91/630/EEC laying down minimum standards for the
1302 protection of pigs. *Official Journal L316*, 01/12/2001,

1303 The Council of The European Union 2001b. Council Directive 2001/88/EC of 23rd October
1304 2001 amending Directive 91/630/EEC laying down minimum standards for the protection of
1305 pigs. *Official Journal L 316*, 01/12/2001,

1306 Trickett SL, Guy JH and Edwards SA 2009. The role of novelty in environmental enrichment
1307 for the weaned pig. *Applied Animal Behaviour Science* 116, 45-51.

1308 Turner SP 2011. Breeding against harmful social behaviours in pigs and chickens: State of
1309 the art and the way forward. *Applied Animal Behaviour Science* 134, 1-9.

1310 Tuytens FAM 2005. The importance of straw for pig and cattle welfare: A review. *Applied*
1311 *Animal Behaviour Science* 92, 261-282.

1312 Valros A, Ahlstrom S, Rintala H, Hakkinen T and Saloniemi H 2004. The prevalence of tail
1313 damage in slaughter pigs in Finland and associations to carcass condemnations. *Acta*
1314 *Agriculturae Scandinavica Section A-Animal Science* 54, 213-219.

1315 Valros A, Palander P, Heinonen M, Brunberg E, Keeling L and Piepponen P 2013.
1316 Preliminary evidence of an altered serotonin metabolism in the prefrontal cortex of tail biting
1317 pigs. ISAE 47th Congress. Florianopolis, Brazil, pg 105.

1318 Van de Perre V, Driessen B, Van Thielen J, Verbeke G and Geers R 2011. Comparison of
1319 pig behaviour when given a sequence of enrichment objects or a chain continuously. *Animal*
1320 *Welfare* 20, 641-649.

1321 Van de Weerd HA and Day JEL 2009. A review of environmental enrichment for pigs housed
1322 in intensive housing systems. *Applied Animal Behaviour Science* 116, 1-20.

1323 Van de Weerd HA, Docking CM, Day JEL, Avery PJ and Edwards SA 2003. A systematic
1324 approach towards developing environmental enrichment for pigs. *Applied Animal Behaviour*
1325 *Science* 84, 101-118.

1326 Van de Weerd HA, Docking CM, Day JEL, Breuer K and Edwards SA 2006. Effects of
1327 species-relevant environmental enrichment on the behaviour and productivity of finishing
1328 pigs. *Applied Animal Behaviour Science* 99, 230-247.

1329 Van de Weerd HA, Docking CM, Day JEL and Edwards SA 2005. The development of
1330 harmful social behaviour in pigs with intact tails and different enrichment backgrounds in two
1331 housing systems. *Animal Science* 80, 289-298.

1332 Van Putten G 1969. An investigation into tail-biting among fattening pigs. *British Veterinary*
1333 *Journal* 125, 511-517.

1334 Verdoes, N 29-1-2014. Starplus stalconcept voor varkens (Starplus Stable concept for pigs)
1335 from <http://www.wageningenur.nl/nl/show/Starplus-stalconcept-voor-varkens.htm>.

1336 Wallenbeck A and Keeling LJ 2013. Using data from electronic feeders on visit frequency
1337 and feed consumption to indicate tail biting outbreaks in commercial pig production. *Journal*
1338 *of Animal Science*.

1339 Wallgren P and Lindahl E 1996. The influence of tail biting on performance of fattening pigs.
1340 *Acta veterinaria scandinavica* 37, 453-460.

1341 Webster SD and Day JEL 1998. Tail-biting and tail-docking in pigs. *The Veterinary Record*
1342 142, 375.

1343 Wechsler B 2013. Experience from Switzerland on ensuring the welfare of pigs. Extended
1344 working group meeting on the development of guidelines concerning Council Directive
1345 2008/120/EC on the protection of pigs (DG SANCO). Brussels, EU.

1346 Westin, R 2003. Svansbitning hos gris relaterat till individuell tillväxt och ras (Tail biting in
1347 pigs related to individual growth and breed). Department of Animal Environment and Health,
1348 Sveriges Lantbruksuniversitet (SLU),
1349 Veterinärmedicinska fakulteten, Veterinärprogrammet,

1350 Wilson K, Zanella R, Ventura C, Johansen HL, Framstad T, Janczak A, Zanella AJ and
1351 Neibergs HL 2012. Identification of chromosomal locations associated with tail biting and
1352 being a victim of tail-biting behaviour in the domestic pig (*Sus scrofa domesticus*). *Journal of*
1353 *Applied Genetics* 53, 449-456.

1354 Zonderland, J.J. 2010. Talking Tails: Quantifying the Development of Tail Biting in Pigs. PhD
1355 Wageningen UR.

1356 Zonderland JJ, Bracke MBM, den Hartog LA, Kemp B and Spoolder HAM 2010a. Gender
1357 effects on tail damage development in single- or mixed-sex groups of weaned piglets.
1358 *Livestock Science* 129, 151-158.

1359 Zonderland JJ, Kemp B, Bracke MBM, den Hartog LA and Spoolder HAM 2010b. Individual
1360 piglets' contribution to the development of tail biting. *Animal* 5, 601-607.

1361 Zonderland JJ, Schepers F, Bracke MBM, den Hartog LA, Kemp B and Spoolder HAM 2011.
1362 Characteristics of biter and victim piglets apparent before a tail-biting outbreak. *Animal* 5,
1363 767-775.

1364 Zonderland JJ, van Riel JW, Bracke MBM, Kemp B, den Hartog LA and Spoolder HAM
1365 2009. Tail posture predicts tail damage among weaned piglets. *Applied Animal Behaviour*
1366 *Science* 121, 165-170.

1367 Zonderland JJ, Wolthuis-Fillerup M, Van Reenen CG, Bracke MBM, Kemp B, den Hartog LA
1368 and Spoolder HAM 2008. Prevention and treatment of tail biting in weaned piglets. *Applied*
1369 *Animal Behaviour Science* 110, 269-281.

1370 Zwicker B, Gygax L, Wechsler B and Weber R 2012. Influence of the accessibility of straw in
1371 racks on exploratory behaviour in finishing pigs. *Livestock Science* 148, 67-73.

1372 Zwicker B, Gygax L, Wechsler B and Weber R 2013. Short- and long-term effects of eight
1373 enrichment materials on the behaviour of finishing pigs fed ad libitum or restrictively. *Applied*
1374 *Animal Behaviour Science* 144, 31-38.
1375
1376

Table 1. Summary of comparative manipulable material studies in which tail injuries were reported for growing pigs. An asterisk (*) by the reference indicates that pigs had intact (undocked) tails, a dagger (†) indicates they were tail docked, and a double dagger (‡) indicates that half the pigs were docked in a 2 x 2 experimental design. All of the studies were controlled experiments with the exception of Courboulayet *al* (2009) which scored tails in part- or fully-slatted and straw systems in an on-farm observational study of 82 farms. Fold-improvement is a result of the tail biting value for the first manipulable material divided by the value for the second. Where two similar materials are named, the average was taken.

Manipulable material A (g/pig/day)	Outcome variable	Value of outcome variable for material A	Manipulable material B (g/pig/day)	Value of outcome variable for material B	Number of pigs (pens) per treatment	P value of difference ⁵	Fold-improvement in outcome of material B over A ³	Mean fold-improvement of similar studies	Log ₁₀ mean fold change	References
None ¹	% removed for tail injury	17.5	Compost rack (500)	<1	108 (6)	<0.05	17.5+	17.5+	1.24	*Beattie <i>et al</i> (2001)
None	% pigs with tail wounds	2.3	Straw bedding	0.5	~4,550 (40 farms)	<0.001	4.6	8.6	0.93	†Courboulayet <i>al</i> (2009)
None	% removed for tail injury	2.5	Straw (490)	0.2	512 (16)	N.S.	12.5			†Scott <i>et al</i> (2007)
Hanging toy	% pigs with tail wounds	42.2	Straw (400)	0	181 (12)	N.A.	(Max)	6.7+	0.83	*Van de Weerd (2005)
Hanging toy	% died or were removed for tail injury	11.7	Straw (400)	1.4	2048 (64)	<0.001	8.4			†Scott <i>et al</i> (2006)
Hanging toy	% pens with tail wounds % removed for tail injury	83 11.1	Straw (5 cm deep)	17 0	72 (6)	P<0.05 N.A.	4.9 (Max)			*Van de Weerd <i>et al</i> (2006)
None	Tail lesion index	0.7	Light chopped straw/wood shavings (12.5)	0.1	126 (31)	P<0.05	7	7	0.85	*Munsterhjelmet <i>al</i> (2009)

Rubber hose or Chain	% of pens with tail wounds	56	Light straw (20)	8	240 (24)	P<0.05	7	7	0.85	*Zonderland <i>et al</i> (2008)
Rubber hose or Chain	% of pens with tail wounds	56	Straw rack (5)	29	240 (24)	N.S./ P<0.05 ²	1.9			*Zonderland <i>et al</i> (2008)
Chain & rubber-covered chain	% prevalence of tail lesions	10.6	Straw rack	12.9	336 (12)	N.S.	0.8			[‡] Scollo <i>et al</i> (2013)
Hanging toy	% pens with tail wounds	83	Straw rack ⁴	50	72 (6)	(N.S.) N.A.	1.7 (7.9)	1.3	0.11	*Van de Weerd <i>et al</i> (2006)
	% removed for tail injury	11.1		1.4						
Rootable feed dispenser	% pens with tail wounds	33	Straw rack ⁴	50	72 (6)	(N.S.) N.A.	0.6 (1)			*Van de Weerd <i>et al</i> (2006)
	% removed for tail injury	1.4		1.4						
Straw rack (5)	% of pens with tail wounds	29	Light straw (20)	8	240 (24)	N.S.	3.6	3.6	0.55	*Zonderland <i>et al</i> (2008)
Straw rack ⁴	% pens with tail wounds	50	Straw (5 cm deep)	17	72 (6)	(N.S.) N.A.	2.9 (Max)	2.9	0.46	*Van de Weerd <i>et al</i> (2006)
	% removed for tail injury	1.4		0						

Footnotes: ¹For the Beattie *et al* (2001) study, 'None' includes the average of pens with nothing and pens with the non-manipulable empty overhead racks.²In this study, straw rack and metal chain had significantly different % tail wounds, but straw rack and rubber hose did not. Metal chain and rubber hose had very similar levels of tail wounds so were combined for simplicity.³'Max' indicates that the improvement was such that tail biting reduced to zero in the second treatment, and this information was not used for the calculation of average fold improvement. Values in parentheses in this column were not used for calculation of the 'Mean fold improvement'- where two different outcome variables were reported for the same study, one of them had to be chosen for use with other studies- the most comparable outcome variables were used where possible.⁴The straw rack was described as a metal tube with a chain mail base which was filled with long straw (and with a tray on the floor underneath) but the quantity provided/used was not reported. ⁵p values are reported where these are available in the source paper. N.S. means that the difference was not

significant, but numerical values have still been used to contribute to estimate the mean fold-change. N.A. means that the p value is not available as it was not reported in the source paper.

Table 2 Comparison of minimum standards for housing grower-finisher pigs across countries and selected assurance schemes (from UK and Denmark) that restrict or completely ban tail docking, with housing standards where docking is widespread (EU, Denmark and UK standard indoor housing).

Country	EU Directives	Denmark	Denmark	Denmark	UK	UK	UK	Sweden	Finland	Norway	Switzerland
System	-	Standard Indoor ³	Antoniussen	Outdoor (includes Organic) ²⁰	Standard Indoor	Freedom Food	Organic ¹⁰	-	-	-	-
Farm Size (finish pigs)	234 ¹	2538 ¹	-	-	1038 ¹	-	-	1046 ¹	485 ¹	264 ¹	420 ¹
Space Allowance 41kg pigs (m2/pig)	0.4 ²	0.4 ³	0.5 ⁴	1.4 (includes outdoor area) ⁵	0.4 ⁷	0.4 (1.17 for straw yard, mucked out monthly) ⁹	555 (0.8 indoor only in extreme weather) ¹⁰	0.48 ¹¹	0.6 ¹³ (0.4 ²¹)	0.5 ¹⁷	0.6 ¹⁹
Space Allowance 101kg pigs (m2/pig)	0.65 ²	0.65 ³	0.85 ⁴	2.3 (includes outdoor area) ⁵	0.65 ⁷	0.75 (1.54 for straw yard, mucked out monthly) ⁹	625 (1.3 indoor only in extreme weather) ¹⁰	0.94 ¹¹	0.9 ¹³ (0.65 ²¹)	0.8 ¹⁷	0.9 ¹⁹
Floor-minimum solid area (% of pen)	0 ²	33 (grower), 50 (weaner) solid or drained by July 2015. Most already comply ³	33-50 ⁴	50 (of indoor area) ⁵	0 ⁷	66 ⁹	50 ¹⁰	70 – 75 ¹¹	67 can include drained floor where perforations are up to 10% of the area ¹³	Solid-floored area large enough for all pigs to lie. ¹⁷	67 lying area, permitted to have 'low degree of perforation for the drainage of liquids' must be solid by 2018 ¹⁹
Tail docking	Not allowed routinely, only if evidence of injuries to ears or tails. "Before..(tail docking).. other measures shall be taken to prevent tail	As EU, but no more than half the tail, and only 2-4 day old piglets) ³ , Docking is	No, but vet can give a time-limited dispensation for	No, but possible to get a dispensation for 60 days for tail	As EU. Docking is widespread. ⁷	Outdoor no, indoor no but can apply for permission to dock to 6cm for 1 yr if they have tail biting (in 2010, 30% did). Must	No ¹⁰	No ¹¹	No ¹⁴	Only by a vet using anaesthetic and long-lasting analgesic ¹⁷	No ¹⁹

	biting and other vices taking into account environment and stocking densities. For this reason inadequate environmental conditions or management systems must be changed." (vet or competent person can dock <7 day old piglets) ²	widespread	r tail biting problems ⁴	biting problems ⁵	take other steps to reduce tail biting to prove docking is a last resort ⁹						
Manip- ulable materials	"Pigs must have permanent access to a sufficient quantity of material to enable proper investigation and manipulation activities, such as straw, hay, wood, sawdust, mushroom compost, peat or a mixture of such, which does not compromise the health of the animals." ²	Denmark wide: Material must be of 'natural origin' and be used 'for rooting' and provided 'on the floor' ³	Straw bedding- all pigs able to lie on straw. ⁴	Straw-bedded indoor lying area, outdoor run can be concrete ⁵	As EU: Chains alone not enough, tyres not allowed, objects not fouled and within reach of pigs ⁷	Lying area and must have comfortable absorbent bedding (straw, sawdust, shredded paper). Permanent access to materials (straw, peat, silages, mushroom compost) in sufficient quantities to allow and encourage proper expression of rooting, pawing and chewing behaviours. ⁹	Mainly outdoor, with soil, stones, green plants. If indoor, 'ample' bedding (straw, sawdust, sand, paper or natural materials such as bracken or rushes, not peat) ¹⁰	Straw must be provided for all pigs ¹¹	Permanent and enough to make into small piles, or if not permanent, materials that can be re-shaped, replenished twice daily (typically straw, sawdust, wood shavings or peat are used), plus additional materials- ball, ¹⁵ chain or sticks. ¹⁵	As EU, but 'wood(chips)' in the list of materials rather than wood ¹⁷	Solid floor bedded with sawdust, straw rack provided ¹⁹
Abattoir scoring to estimate tail biting prevalence		0.5 - 1.5% ⁶	-	1.0-4.0% ⁶	1.0% 'severe tail lesions' 2.4% 'evidence of tail biting' ⁸	-	-	<2.0% tail damage ¹²	1.8% tail damage, 5.1% partial condemnations ¹⁶	4.0% tail damage ¹⁸	

Footnotes: 1) Farm sizes were calculated from Eurostat (2013) figures for 2010 for 'other pigs' which includes grower/finisher pigs (available only at country level). Low EU average is due to the inclusion of many member states which do not have a major pig industry. 2) EU Directives 2001/88/EC and 2001/93/EC (The Council of The European Union, 2001a; The Council of The European Union, 2001b). 3) Ban on fully-slatted floors applies from July 2000 for new buildings, and for all housing by July 2015. Drained floor defined as maximum 10% openings. Danish Government (2000; 2003a; 2003b) DVFA (2013) 4) Antonius: Danish Crown (2007) 5) Outdoor: Friland (2012), Ministeriet for Fødevarer, Landbrug og Fiskeri (MFLF, 2012) 6) Taken from figure j, p86 in Forkman *et al.* (2010) 2008-2010 figures from one abattoir so are directly comparable between systems. 7) Defra (2003), BPEX (2010) 1% figure from Northern Ireland and Republic of Ireland (Harley *et al.*, 2012), 2.4% for 6 abattoirs in England (Hunter *et al.*, 2001) 9) RSPCA (2012) and Kate Parkes (RSPCA pers. comm.) 10) Soil Association (2012), Outdoor-based system, giving permanent access to soil and growing plant foods. Must provide summer wallows and/or shade. Rotational grazing required. Indoors only under exceptional circumstances and must have outside run allowing rooting and dunging. 11) Jordbruksverket (2010), Mulet *et al.* (2010) SLU and LRF (2009), 12) Holmgren & Lundeheim (2004), Keeling *et al.* (2012). 13) Council of State (2012). The regulations came into effect in the first of January 2013. If a facility was already operating at that time, the space allowance regulations come into effect on the first of January 2018, and the minimum solid floor area on the first of January 2028, or both come into effect upon renovation if that is sooner. 14) Tail docking prohibited since January 2003 (Council of State, 2002). 15) Evira (2013) 16) Partanen *et al.* (2012) 17) LMD (2003) 18) Fjetland & Kjastad (2002) 19) Swiss Federal Council (2008), Wechsler (2013), CIWF (2009) 20) Outdoor access (can be concrete): 0.6m²/pig at 40kg and 1.0m²/pig at 100kg. 21) MAF (1997). Applies to all facilities until the 31st of December 2012 and to old facilities not renovated before 2018

Figure 1: Postulated relationships between the underlying processes of tail biting (text in bold, connected in order by solid arrows) and various known or suspected risk factors (text in plain type) connected with 19 dashed numbered arrows to show how some of the risk factors might influence each other or the underlying process of tail biting. Some proposed risk factors for which the evidence is currently weak (e.g. disease and parasitism, draughts) are included where a plausible hypothesis exists. The meaning of the numbered arrows is explained in Supplementary Material S1.

Figure 2: Enrichment materials' relative effect at reducing tail biting based on Log₁₀ fold reductions in tail damage, using studies from Table 1.

Footnotes:

Line thickness indicates the number of studies used; thinnest lines = 1 study, intermediate lines = 2 studies, thickest lines = 3 studies. Shading of the box indicates the amount of material that is used up. Compost and Straw (shown in black), at least 500 g/pig/day, light straw (in dark grey, 12.5 to 20 g/pig/day), straw rack (5 g/pig/day). None and Straw used as reference, as these are the most common materials used across studies. This means that none and straw each have only one horizontal line. Light straw and Straw rack have multiple lines, which show the range of positions they could occupy relative to other substrates based on a number of studies.